



MIKE WATKINS, MPA, RN,FP-C
VIRGINIA COMMONWEALTH
UNIVERSITY HEALTH SYSTEM



Case Concepts of Capnography

Objectives

- Brief Review Capnography
 - History
 - Anatomy and Physiology
- Case Presentations
 - Pathophysiology of Respiration and Ventilation
 - Capnography integrated with the critical care patient
 - “Advanced” application of capnography
 - Alternative Applications

So you have
them intubated!

Next step??



OR ...

**YOU ARE TAKING OVER
SOMEONE ELSE'S
AIRWAY**

First Priority

- Maintenance of Intubation
 - Confirm and re-confirm placement
 - Visualization
 - Auscultation
 - Observation
- Capnography:
 - “Memorial, we have confirmed tube placement with a BEAUTIFUL BOX SHAPED waveform at 35 to 40!”

Confirmation

- When you put the tube in the trachea
 - Watching it pass through the cords
- When your assistant moves the BVM the wrong way
 - When the hairs on your neck stand up
- When you move the patient
 - From the house, to the stretcher, to the ambulance, to the hospital, on the code bed

Capnography

- Quantitative, graphical measurement of instantaneous CO₂ concentration
 - American Society of Anesthesiologists (ASA) standards:
 - Every patient receiving anesthesia shall have adequacy of ventilation continuously monitored
 - Continual monitoring for the presence of expired carbon dioxide shall be performed unless invalidated by the nature of patient, procedure, or equipment
 - Continual EtCO₂ analysis, in use from the time of ET placement, until extubation (removal of the trachea), shall be performed using a quantitative method such as infrared spectrometry, or mass spectroscopy
- EVERY INTUBATED PATIENT..**
Continuous Monitoring...
Quantitative.

History of Capnography

- Developed in 1961 (For practical purposes)
- Expensive and bulky product limited to OR anesthesia
- In mid 1980s, anesthesia related fatalities led to need for improvements in airway management
 - Malpractice costs rise 1975 to 1985

History of Capnography

- 1988: Anesthesia standardized use of pulse oximeters and capnography
 - Malpractice claims from hypoxic related injury almost eliminated -Massachusetts
 - Insurance claims from anesthesia drops from 11% to 3% over 15 years
 - In 2002, anesthesia insurance premium was \$18,000, the same as it was in 1985

Integration of Capnography

- In Anesthesia, capnography is an industry standard
- In EMS, it is a standard, but not there are variables
 - "No, its not the pulse ox!"
 - Waveform versus colorimetric
- It is not common in Emergency Departments, and varied in ICUs


Patient Safety

ABOVE ALL, DO NO HARM

A&P for Capnography:

- What is important:
 - Air movement
 - Surface area of lungs
 - Blood flow to lungs and body
- Respiratory Cycle
 - Alveolar level
 - Cellular level

Respiratory Anatomy



Review of Metabolism

- **Aerobic:**
 - Oxygen and Glucose metabolize to produce Energy to do work
 - Carbon Dioxide and Water are the byproducts
 - Krebs Cycle
 - Most efficient process
 - Improves with exercise



Review of Metabolism

- **Anerobic:**
 - Lack of oxygen causes build up of acids
 - Lactic Acid and Pyruvic Acid
- **Buffer System**
 - Hydrogen Ions of the Acid (pH) combine with Bicarbonate to form Carbonic Acid
 - This breaks down into water and carbon dioxide
 - Increased CO₂ stimulate increased ventilation rate to remove it

Carbon Dioxide

- By-product of normal respiration
- Measured as a Partial Pressure
 - 35-45 Mm/Hg
- Measured as a Percentage
 - 5-6%
- Key for: respiratory drive, ph balance
- Considered “acidic”

Drive to Breathe

- CO₂ triggers breathing
- **NOT ENOUGH**
 - Hypoventilation leads to hypercarbia
 - Hypercarbia leads to respiratory acidosis
- **TOO MUCH**
 - Hyperventilation leads to hypocarbia
 - Hypocarbia leads to respiratory alkalosis

What do the numbers mean?

- **Oxygen and Carbon Dioxide**
- **Hypoventilation:**
 - O₂ < 60mm/Hg
 - CO₂ > 45mm/Hg (Hypercapnea)
- **Hyperventilation:**
 - O₂ > 100mm/hg (SaO₂ above 98%)
 - CO₂ < 35mm/Hg

CO₂ on the BRAIN

- Decreased CO₂ from hyperventilation
 - Cerebral Vasoconstriction
 - Balancing
- Indication: (old school)
 - Traumatic head injury/CVA
 - Maintain perfusion without worsening bleeding
 - End-tidal CO₂ target is 33 to 35mm/Hg

CO₂ on the Brain

- Elevated CO₂
 - Permissive Hypercarbia
 - Above 45mm/Hg
 - With adequate oxygenation
- Potential Benefits
 - Cerebral and systemic vasodilation
 - Increase cellular oxygen supply
 - Decrease oxygen demand

Priority is Oxygenation

- Adequate Oxygenation
- Adequate Ventilation
- Avoid tunnel vision
 - Use your tools
- Quality over Quantity

HYPOXIA KILLS

Dalton's Law: Partial Pressure of Gas

Total pressure of a gas is equal to the SUM of the partial pressures of the gas


- Atmospheric pressure is 760mm/Hg at sea level
- Under NORMAL conditions, all of the atmospheric gas pressures add up to 760
 - Oxygen is 159.2 mm/Hg
 - Nitrogen is 592.8 mm/Hg
 - CO₂ is 0.23 mm/Hg
 - Other gases, like Argon = 8mm/Hg

Atmospheric Gases

- Convert percentage to pressure
- Normal gas Percentage
 - Oxygen at sea level: 21%
 - CO₂ and other gases: 1%
 - Nitrogen: 78%

Partial Pressure

- Gradient
- The exchange of gases based on pressure gradient
 - Pressure forces Oxygen onto Hemoglobin




Ventilation and perfusion

Pathology that Impacts CO₂

- **Ventilation Problems**
 - Inability to move air in and out of the alveoli
 - Hyperventilation, hypoventilation
- **Perfusion Problems**
 - Oxygen transport to cells
 - Lack of blood flow
 - Ability of blood to carry oxygen

Normal Ventilation/Perfusion



Alveolar Perfusion Problems

- **Shunt Problem**
 - Blocking of bronchial airways
 - Pneumonia, atelectasis
 - Right mainstem intubation
 - Causes retention of CO₂, increased levels
- **Dead Space Ventilation**
 - Capillary flow to alveoli impaired
 - Low Cardiac output, hypotension
 - Excessive PEEP
 - CO₂ does not cross into the alveoli for exhalation
 - Decreased levels of CO₂


Clinical Conditions with Increased CO₂

- **Increased CO₂ production**
 - *Bicarbonate administration, fever, seizures, sepsis, thyroid storm*
- **Decreased alveolar ventilation**
 - *Hypoventilation, muscular paralysis, respiratory depression, COPD (retaining CO₂)*
- **Equipment Problem**
 - *Rebreathing, ventilator leak*

Clinical Conditions: Decreased CO₂

- **Decreased CO₂ production**
 - *Cardiac arrest, hypotension, hypothermia, pulmonary emboli, pulmonary hypoperfusion*
- **Increased alveolar ventilation**
 - *Hyperventilation*
- **Equipment Problems**
 - *Airway obstruction, esophageal intubation, ETT leak, incomplete exhalation, poor sampling, ventilator disconnect*

Normal EtCO₂ waveform



Troubleshooting

- False Positive
 - Large amount of carbonated beverage
 - AHA
 - Rapidly Declines
- False Negative
 - More common
 - Low flow states
 - Air movement
 - Blood Flow

Troubleshooting


- Sudden loss of waveform
 - IMMEDIATE CLINICAL RECONFIRMATION
 - Lung sounds, SaO₂, Anything else
- Place colorimetric detector
- Clean/Clear sensor
 - Blockage
- Recalibrate/zero if able
- Replace adapter

Case Presentations




- What are you seeing?
 - What does your physical assessment tell you?
- What are your transport considerations?
 - Interventions
- Differential Diagnosis?
- Trouble shooting?
 - Is the data valid?

Case 1

- Respiratory Distress
- 54 y/o COPD
- Respiratory Rate: 24
- Pursed lips



- ### Obstructive Airway Disease
- Shunt problem
 - Asthma, COPD, Emphysema
 - Swelling of airways/excess mucus
 - Airflow turbulent
 - Forceful expiration
 - Different EtCO₂ presentations:
 - Mild=hyperventilation, low EtCO₂
 - Moderate=normal EtCO₂, waveform change
 - Severe=elevated EtCO₂, sharkfin




- ### Case 2
- 65 year old obese trauma patient
 - Predicted Difficult Intubation
 - Multiple Injuries
 - Chest Contusions
 - Abdominal Distention
 - Fractures of right upper leg, left lower leg, and right arm
 - Complains of Respiratory Distress

Case 2

- Initial Et CO₂ 6-7mm/Hg
- Intermittent sensor detection of numerical value
- Waveform present
- Low “shark fin” appearance
- What is going on?
- Is the ET good?

Case 2



Shock

- “A rude unhooking of the Machinery of Life”
 - Samuel Gross, 1872
- “A momentary pause in the act of death”
 - John Collins Warren, 1895
- “Pushing back the edge of death”
 - Judy Mikhail, 1999

Shock


- As the body’s compensatory mechanisms begin to work, we appreciate changes in vitals signs:
 - Altered mental status
 - Pale, clammy, diaphoretic
 - Increased heart rate and respiration
 - Decreased blood pressure

Shock and Capnography

- A Piece of the Puzzle
- Anaerobic compensation for decreased perfusion:
 - CO₂ elevated, ventilations increase
- Cardiac Output drops:
 - Vasodilation or from hypovolemia:
 - CO₂ decreased as detected by EtCO₂




Shock

- Capnography:
 - Index of Resuscitation
 - Quality of perfusion
 - Quality of ventilation
- Like ALL monitors, it is a TOOL
 - Understand its limitations
 - It can GUIDE decisions
 - It should not MAKE the decision




Case 3

- 26 year old MVC at community hospital
- Intubated in ED after becoming combative
- Vitals: Respirations 20 assisted on ventilator
- Assist/Control:
 - Rate12, TV 500, FiO₂ 50%, PSV 10, PEEP 5




Case 4

- 50 year old cancer patient receiving radiation and chemo
- Presents with respiratory distress to ED
 - SaO₂: 85%, dramatic work of breathing, becoming tired
 - intubated promptly, placed on ventilator
 - Vitals: BP 140/88, HR 78, vented at 10 with SaO₂ of 93% with 100% FiO₂
 - Initial EtCO₂ is 20mmHg




- ### Case 4
- Community ED requests transfer to tertiary care for Pulmonary Embolism
 - Post Intubation ABG:
 - pH 7.31, PaO₂: 140, PaCO₂: 49mmHg, Bicarb 27
 - CO₂ gradient:
 - PaCO₂ – PetCO₂ (49 minus 20 equals 29mmHg)
 - Normal gradient 3 to 5mmHg
 - What is in the blood is not getting out

- ### Pulmonary Embolism
- Dead Space Ventilation
 - Decreased EtCO₂
 - Clot breaks loose in blood vessel
 - Floats to and obstructs pulmonary vasculature
 - Causes:
 - Post surgical
 - Sitting for extended time
 - David Bloom, NBC News in 2003



- ### ABGs, pH, and Capnography
- Arterial Blood Gases assess for acid-base balance
 - Acidosis and Alkalosis
 - Mechanisms: Respiratory and Renal (Metabolic)
 - pH is a measure of Hydrogen ion concentration (H⁺)
 - Normal is 7.35 to 7.45
 - Reflects balance between carbon dioxide and bicarbonate
 - Capnography only represents the RESPIRATORY




Case 5


- 21 year old female
- Witnesses cardiac arrest on athletic track, defibrillated by AED
- BLS and ALS procedures per protocol
- No pulses or vitals



Case 5 Initial: No pulses




Case 5: No pulses



Cardiac Arrest and CO2


- In cardiopulmonary arrest
 - CO2 levels in blood stream increase
 - Exhaled CO2 levels decrease due to low flow states
 - No ventilation, no circulation of blood
 - No perfusion
 - Cambridge journal Article
- Return of Spontaneous Circulation (ROSC)
 - Spike in EtCO2 after trend of low levels

Case 5 Variant





Case 6

- 48 year old COPD
- Cyanotic, lethargic
- Vitals: HR: 131 A-fib, BP: 158/100, RR: 32, SaO2 on NRB: 90%
- Previous Intubations for same
- Airway Considerations?
- Attempt Intubation...




- Case 7**
- 2 year old female with new onset seizures
 - Inter-facility transport for tertiary care
 - Intubated / Ventilated by BVM
 - 4.5 ET uncuffed
 - Vitals: HR 160, BP 84/40, Ventilated at rate of 36, SaO₂ 100%, Temp: 103
 - Ventilator: FiO₂ 100%, PEEP 5, initial I:E 1:2.7

- Case 7**
- Capnography
 - Initial level: 98mm/Hg
 - Shape: elevated box shape, irregular respiratory pattern at rate of 36
 - What are your actions?
 - Increase rate?
 - Change I:E ratio?
 - ET problem
 - How might etiology change treatment?
 - Asthma
 - Trauma



- Case 8**
- Interfacility transport:
 - 56 year old male admitted with "fever"
 - Diagnosed with "sepsis:
 - PMH: ESRD, IDDM, CAD, CHF
 - Lethargic, GCS 12,
 - Vitals: 84/60. HR 130, respirations 10 irregular
 - Intubated electively for transport
 - Initial EtCO₂: 21mmg/Hg

Case 8

CO₂

Case 8

CO₂

- Capnography:
 - Reason for Low EtCO₂?
- What are your corrective actions?
 - Decrease ventilation rate?
 - Fluids?
 - Pressors?
 - Blood products?

Case 8

CO₂

- Consider:
 - BP of 140/90, HR 110, RR 28
- EtCO₂ of 28mHg
- Respiratory alkalosis as an initial compensation for metabolic acidosis
 - Capnography considered a potential triage tool.

Case 9

CO₂

- Male patient with arm trapped in roller press
- Awake, oriented, agitated and in severe pain, 10/10
 - BP 150/70, HR 118, R 20, SaO₂ 100%
- Movement of arm increases agitation and reduces access
- Movement of rollers causes pain
- Elect to sedate for extrication
 - Online medical control

Case 9

CO₂

Case 9

CO₂

- Need for pain control and sedation
- Patient is in difficult position to monitor vitals
- What is your pain control protocol?
- Do you have a sedation protocol?
- How much is too much?
 - Capnography by sidestream


Case 9: Sedation

- Quality of ventilation
- Detection of Apnea
- Predictor of Compromise
- Out-of Hospital sedation:
 - Long distance and air medical transport
 - Extrinsic


Case 10

- 21 year old male c/c chest pains
- Sudden onset
- Stabbing, non-radiating, 10/10
- Tingling in his fingers

Case 10



Variety



Case 10

23:02:16	Initial Oxygen	94	---	23:05	
23:05:25	Vital Signs	83	97-95	23:08	25/77/90-92
23:06:58	HRIP	79	98-99	23:13	
23:12:25	HRIP	83	97-94	23:13	100/60/70/100
23:15:25	HRIP	78	94-94	23:26	
23:26:12	HRIP	73	98-92	23:26	
23:26:38	HRIP	73	98-92	23:25	80/64/70/92

End-tidal by Nasal Prongs




Case 11

- Female trauma patient
 - Unrestrained driver with steering wheel deformity; found under dash after airbag deployed
 - Pattern of injury??
- Intubated successfully
 - Confirmed by waveform sedated and paralyzed
- 25 minute flight to Trauma Center

Case 11: Cruising Along




CO₂



Case 11: Sudden Change!



CO₂



Case 12



- 57 year old obese male with spinal trauma
 - Fell forward, hyper-flexion of neck
 - Confirmed C₅, C₆ fractures
- CNS Intact-full movement
 - GCS of 9T (14 if not intubated)
- Intubated: and we did not know how
 - Sedation/fiberoptic ETI by anesthesia
- On T-piece, NOT VENTILATED breathing on his own

Case 12




- Considerations:
 - Community hospital to Level 1 trauma Center
 - Patient obese: 280 pounds
 - Aircraft: EC135
 - No existing ventilator settings; crew discretion on "optimal"
 - Difficult airway on multiple dimensions
 - Confirmed by CXR prior to movement

Case 12: Volume Ventilation




CO₂



Case 12: Pressure Ventilation



CO₂



Case 12


- Pressure versus volume ventilation
 - Pressure Control 24, FiO₂ 100%, Rate 12, Assist Control with PSV
 - Switch to BVM after desaturation
- Sedation, paralysis, pain control
- How might a different airframe change management? Or ground unit?
 - Bell 407 vs. 412
 - EC130 vs EC 135/145

Case 12

- Physical restriction of breathing
 - Burns
 - COPD
 - Trauma
 - Surgical

Case 13

- 24 y/o patient in head on MVC
- Altered LOC, combative, signs of head injury
- RSI clinical course
 - Etomidate 0.3 mg/kg
 - Succinylcholine 1.5 mg/kg
- After paralytic, patient developed trismus and rigidity
- Unable to intubate, but can ventilate with oral airway in place
 - Unable to open mouth to place King LTD

CO₂

Case 13 Malignant Hyperthermia

- Life Threatening
- Hypermetabolic state in patient's with hereditary skeletal muscle defect
 - Genetic predisposition 1:10000
 - Clinical Incidence 1:30000
- Depolarizing muscle relaxants (Succinylcholine) and anesthetic gases cause raise in myoplasmic calcium

Malignant Hyperthermia: Signs and Symptoms

- Hypercarbia: most sensitive indicator in intubated patient
- Tachycardia
- Tachypnea
- Temperature elevation
- Hypertension
- Dysrhythmias
- Acidosis
- Hypoxia
- Hyperkalemia
- Skeletal muscle rigidity
- Myoglobinuria

MH Management

- Get help: Medical control
- Hyperventilate patient with 100% oxygen
- Cool patient
- Antidote is Dantrolene
 - Truly the only effective treatment
 - Operating rooms have an MH cart stocked with multiple bottles
- Prehospital considerations
 - Non-depolarizing paralytic
 - Benzodiazepines

Critical Care Transport

- Verification of ETI, as with scene
- Evaluation of ventilation
 - Alter ventilator settings
- Evaluation tool for perfusion
- Simple information will help reflect in large changes
 - Decrease ventilation rate
 - Improve quality of chest compressions

Transition Monitoring

- Specific to non-cardiac, trauma patients
- Handheld device in pouch: combine EtCO₂ and SaO₂
- Advantages:
 - Reduces scene and movement times
 - Provides critical but NOT complete information
 - Reduces bulk



Summary:

- Capnography is a TOOL
 - Does not substitute for good clinical skills
- Remember the BASICS
 - ABCs
- DO NOT OVERTHINK Capnography
 - Some cases will be difficult to figure out

Sources:

- Ahrens, T. D., & Sona, C. M. (2003). Capnography Application in Acute and Critical Care. *AACN Clinical Issues*, 14 (2), 123-132.
- American Society of Anesthesiologists. (2005, October 25). *Publications and Services*. Retrieved September 16, 2007, from American Society of Anesthesiologists: <http://www.asahq.org/publicationsAndServices/sgstoc.htm>
- Brain Trauma Foundation. (2000). *Guidelines for Prehospital Treatment of Traumatic Brain Injury*. New York: Brain Trauma Foundation.
- Cooper, J. B. *Medical Technology: Patient Safety is Paramount*.
- Garey, B. (2007, August 18). Flight Paramedic, Medflight I. (M. Watkins, Interviewer)
- Gravenstein, J. S., Jaffe, M. B., & Paulus, D. A. (2004). *Capnography*. Cambridge, United Kingdom: Cambridge University Press.
- Hassett, P., & Laffey, J. G. (2007). Permissive Hypercarbia: Balancing Risks and Benefits in the peripheral microcirculation. *Critical Care Medicine*, 2229-2230.
- Huether, S. E., & McCance, K. L. (2004). *Understanding Pathophysiology* (3 ed.). St. Louis, Missouri: Mosby Elsevier.
- Hutchison, R., & Rodriguez, L. (2008). Capnography and Respiratory Depression. *American Journal of Nursing*, 108 (2), 35-39.
- McCillcuddy, D., Tang, A., Cataldo, L., Guere, J., & Shapiro, N. I. (2008). Evaluation of end-tidal carbon dioxide role in predicting elevated. *Intern Emergency Medicine*. pre-pubish.



THANK YOU!!!!

ANY QUESTIONS?

