

Purpose of Form

The TB Risk Form (TB 512) is a tool to assess and document a patient's symptoms and/or risk factors. Completing this form will also help in determining the need for future medical testing and evaluation.

Directions for Completing the Form

Print clearly and complete this form according to the instructions provided below.

I. Screen for Presence of TB Symptoms

Screen the patient for symptoms of active TB disease

- All symptomatic individuals should: (1) receive a test for TB infection if not previously positive (TST or IGRA); (2) have their sputum collected; (3) be referred for an immediate chest x-ray and medical evaluation, regardless of the TST result.
- If the patient does not have symptoms of active TB disease, go to Section II and assess risk for LTBI and/or disease.
- *Symptoms of active TB disease are more subtle in children.* Children with symptoms of active TB disease should receive a TST, CXR and immediate medical evaluation by medical personnel knowledgeable about pediatric TB.

II. Screen for TB Infection Risk (In subsections A and B, check all the risk factors that apply)

Section II has 2 sections. Section A: "Assess Risk for Acquiring LTBI", Section B: "Assess Risk for Developing TB Disease if infected".

- If a patient has one or more risk factors for LTBI as listed in sections A or B, then go to Section III and administer the TST or IGRA.
- If a patient does not have risk factors for LTBI, do not administer the TST or IGRA. Go to Section III and place a check next to "No Risk Factors for TB Infection."
- If the patient's school, employment, etc. requires a TB screening, place a check next "Issued Screening Letter" (Section IV) and provide this document to the patient.

A. Assess Risk for Acquiring LTBI -- The following are definitions of select categories of persons at risk for LTBI

B. Assess Risk for Developing TB Disease if infected - The following are definitions of select categories of persons at risk for TB disease if infected

- *Person is a current close contact of another individual known or suspected to have TB disease --*
Person is part of a current TB contact investigation
- *Lived in or visited another country where TB is common for 3 months or more, regardless of time in the U.S. --*
Person lived or visited a high endemic country ≥ 3 months. High endemic country is defined as a case rate of $\geq 20/100,000$. See VDH Exception list for LOW risk countries.
- *Person is a resident/employee of high TB risk congregate settings--*
These settings are correctional facilities, nursing homes, and long-term care institutions for the elderly, mentally ill and persons with AIDS.
- *Person is a health care worker who serves high risk clients --*
Screen for the individual risk factors for TB infection, unless screening efforts are part of an ongoing facility infection control program approved by local health department.
- *Person is medically underserved --*
Person doesn't have a regular health care provider, and has not received medical care within the last 2 years.
- *Person is an infant, a child or an adolescent exposed to an adult(s) in high-risk categories --*
Child has foreign-born parents, or child's parents/caretakers are at high risk for acquiring TB infection.
- *Person is a member of a group identified by a local health department to be at an increased risk for TB infection --*
Identification of a group is based on local epidemiologic data showing an increase in the number of persons with TB disease or TB infection in the given group.
- *Person needs baseline/annual testing approved by health department --* includes those entering health professions; new health care workers need 2-step TST unless documented negative TST in prior 12 months. A single IGRA is also acceptable. May include screening program that is approved by the local health dept. for facilities or individuals at an increased risk for LTBI.

- *Person's HIV Status is unknown but has risk for HIV infection--*
Offer HIV test. Administer the TB Skin Test, even if the patient refuses the HIV test.
- *Person with clinical conditions that place them at high risk --*
Conditions include substance abuse, chest x-ray findings that suggest previous TB, diabetes mellitus, silicosis, prolonged corticosteroid therapy, cancer of the head and neck, leukemia, lymphoma, hematologic and reticuloendothelial diseases, end stage renal disease, intestinal bypass or gastrectomy, and chronic malabsorption syndromes.
- *Person is on immunosuppressive therapy --*
Person is taking ≥ 15 mg/day of prednisone for ≥ 1 month; person is receiving treatment with TNF- α antagonists (Remicad, Humira, etc.) or other biologic response modifiers and/or person needs baseline evaluation prior to start of treatment with the medications cited here.

III. Finding(s) (Check all findings that apply.)

In this section, indicate findings from the assessments in all previous sections.

IV. Action(s) (Check all actions that apply.)

- Indicate the action(s) to take as a result of the findings in Section III.
- If administering a TB Skin test or IGRA, provide all requested data.
- Repeat TB Skin test or IGRA if appropriate.

Additional follow-up to a Mantoux TB skin test or IGRA blood test

- If the patient's TST reaction or IGRA is interpreted as positive or if she/he has symptoms for TB disease, refer the patient immediately for medical evaluation and a chest x-ray.
- If a person has a history of a positive TST or IGRA and is currently asymptomatic, then refer for a chest x-ray if the following two conditions apply: 1) patient is a candidate for LTBI treatment; **and**, 2) patient is willing to adhere to the treatment.
- If treatment for LTBI is not planned and TB was previously ruled out with a normal chest x-ray, then repeat chest x-rays are not indicated unless symptomatic.