

Table 4: Number (%) of Cases* Reported by Condition, Region of Residence and MMWR Month/Year, Virginia 2016

Condition	January-November 2016										
	Central		Eastern		Northern		Northwest		Southwest		Total
	Cases	%	Cases	%	Cases	%	Cases	%	Cases	%	
Amebiasis	4	12.1%	2	6.1%	22	66.7%	3	9.1%	2	6.1%	33
Botulism	1	20.0%	1	20.0%	0	0.0%	3	60.0%	0	0.0%	5
Brucellosis	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0
Campylobacteriosis	171	12.2%	193	13.7%	486	34.6%	396	28.2%	159	11.3%	1,405
Chikungunya	0	0.0%	0	0.0%	4	80.0%	0	0.0%	1	20.0%	5
Cryptosporidiosis	12	5.6%	68	31.5%	71	32.9%	44	20.4%	21	9.7%	216
Cyclosporiasis	0	0.0%	0	0.0%	7	0.0%	4	0.0%	0	0.0%	11
Dengue	1	4.0%	3	12.0%	17	68.0%	2	8.0%	2	8.0%	25
Ehrlichiosis/Anaplasmosis	25	27.2%	10	10.9%	8	8.7%	23	25.0%	26	28.3%	92
<i>E. coli</i> (STEC)	7	5.9%	5	4.2%	48	40.3%	36	30.3%	23	19.3%	119
Giardiasis	32	11.8%	43	15.8%	131	48.2%	39	14.3%	27	9.9%	272
<i>Haemophilus influenzae</i> , invasive	26	23.2%	24	21.4%	21	18.8%	22	19.6%	19	17.0%	112
Hemolytic uremic syndrome	0	0.0%	0	0.0%	1	33.3%	2	66.7%	0	0.0%	3
Hepatitis A	16	8.6%	27	14.6%	72	38.9%	58	31.4%	12	6.5%	185
Hepatitis B, acute	17	33.3%	9	17.6%	3	5.9%	3	5.9%	19	37.3%	51
Hepatitis B, chronic	186	15.2%	217	17.8%	590	48.4%	111	9.1%	116	9.5%	1,220
Hepatitis C, acute	9	15.0%	3	5.0%	5	8.3%	24	40.0%	19	31.7%	60
Hepatitis C, chronic	1,994	23.5%	1,921	22.6%	1,297	15.3%	1,183	13.9%	2,107	24.8%	8,502
Lead, elevated levels <16 years	176	21.9%	105	13.0%	297	36.9%	107	13.3%	120	14.9%	805
Legionellosis	34	29.6%	31	27.0%	19	16.5%	13	11.3%	18	15.7%	115
Leprosy	0	0.0%	0	0.0%	1	100.0%	0	0.0%	0	0.0%	1
Listeriosis	7	35.0%	2	10.0%	4	20.0%	3	15.0%	4	20.0%	20
Lyme disease	52	4.7%	25	2.3%	349	31.6%	269	24.3%	410	37.1%	1,105
Malaria	2	3.3%	6	9.8%	48	78.7%	5	8.2%	0	0.0%	61
Measles	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0
Meningococcal disease	1	11.1%	0	0.0%	3	33.3%	3	33.3%	2	22.2%	9
Mumps	2	14.3%	0	0.0%	10	71.4%	1	7.1%	1	7.1%	14
Pertussis	19	11.0%	16	9.3%	21	12.2%	68	39.5%	48	27.9%	172
Salmonellosis	187	17.1%	245	22.5%	300	27.5%	203	18.6%	156	14.3%	1,091
Shigellosis	51	15.6%	166	50.9%	50	15.3%	18	5.5%	41	12.6%	326
Spotted fever rickettsiosis	76	27.4%	25	9.0%	34	12.3%	55	19.9%	87	31.4%	277
Streptococcal Group A, invasive	37	18.3%	55	27.2%	40	19.8%	31	15.3%	39	19.3%	202
Streptococcal Group A, toxic shock	0	0.0%	1	25.0%	1	25.0%	0	0.0%	2	50.0%	4
<i>Streptococcus pneumoniae</i> , invasive <5	2	6.7%	5	16.7%	9	30.0%	9	30.0%	5	16.7%	30
Tuberculosis	20	11.8%	26	15.4%	101	59.8%	12	7.1%	10	5.9%	169
Tularemia	2	100.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	2
Typhoid fever	1	8.3%	1	8.3%	8	66.7%	0	0.0%	2	16.7%	12
Varicella (chickenpox)	26	11.0%	22	9.3%	124	52.3%	35	14.8%	30	12.7%	237
Vibriosis	10	25.6%	22	56.4%	3	7.7%	4	10.3%	0	0.0%	39
West Nile virus	2	0.0%	0	0.0%	3	0.0%	3	0.0%	1	0.0%	9
Yersiniosis	1	5.6%	2	11.1%	3	16.7%	6	33.3%	6	33.3%	18

* 2016 data is preliminary, data extracted on 12/13/2016 @ 7:44 am