

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Northern) Middle School Survey Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN6: Among students who rode a bicycle, the percentage who never or rarely wore a bicycle helmet	50.6	1,233	5.59	(38.6 - 62.5)	15.4
QN7: Among students who used rollerblades or rode a skateboard, the percentage who never or rarely wore a helmet	55.5	727	5.71	(43.0 - 67.3)	9.6
QN8: Percentage of students who never or rarely wore a seat belt when riding in a car	3.8	1,461	0.54	(2.8 - 5.2)	1.2
QN9: Percentage of students who ever rode in a car driven by someone who had been drinking alcohol	13.9	1,436	1.60	(10.8 - 17.8)	3.1
QN10: Percentage of students who ever carried a weapon, such as a gun, knife, or club	26.0	1,445	3.36	(19.4 - 34.0)	8.5
QN11: Percentage of students who have ever been in a physical fight	34.3	1,451	2.98	(28.1 - 41.0)	5.7
QN12: Percentage of students who were ever in a physical fight in which they were hurt and had to be treated by a doctor or nurse	3.4	1,455	0.48	(2.5 - 4.7)	1.0
QN13: Percentage of students who had ever been bullied on school property	40.5	1,460	1.88	(36.5 - 44.7)	2.1
QN14: Percentage of students who had ever been electronically bullied	23.0	1,458	2.36	(18.3 - 28.5)	4.6

*The average design effect across all items for this survey is 4.1.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Northern) Middle School Survey Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN15: Percentage of students who ever seriously thought about killing themselves	14.5	1,435	1.59	(11.3 - 18.3)	2.9
QN16: Percentage of students who ever made a plan about how they would kill themselves	10.2	1,458	0.97	(8.3 - 12.5)	1.5
QN17: Percentage of students who ever tried to kill themselves	5.0	1,464	0.65	(3.8 - 6.7)	1.3
QN18: Percentage of students who ever tried cigarette smoking, even one or two puffs	10.4	1,380	2.43	(6.2 - 17.1)	8.7
QN19: Percentage of students who smoked a whole cigarette for the first time before age 11 years	2.5	1,382	0.62	(1.5 - 4.3)	2.1
QN20: Percentage of students who smoked cigarettes on one or more of the past 30 days	1.1	1,392	0.34	(0.6 - 2.1)	1.4
QN21: Among students who reported current cigarette use, the percentage who smoked more than 10 cigarettes per day on the days they smoked during the past 30 days	6.3	15	6.50	(0.6 - 42.6)	1.0
QN22: Among students who reported current cigarette use, the percentage who usually got their own cigarettes by buying them in a store or gas station during the past 30 days	13.3	16	5.95	(4.8 - 32.1)	0.5
QN23: Percentage of students who ever smoked cigarettes daily, that is, at least one cigarette every day for 30 days	0.9	1,415	0.32	(0.4 - 1.9)	1.6

*The average design effect across all items for this survey is 4.1.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Northern) Middle School Survey Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN24: Percentage of students who used chewing tobacco, snuff, or dip on one or more of the past 30 days	1.1	1,464	0.36	(0.6 - 2.2)	1.7
QN25: Percentage of students who smoked cigars, cigarillos, or little cigars on one or more of the past 30 days	0.9	1,464	0.32	(0.4 - 2.0)	1.7
QN26: Percentage of students who ever had a drink of alcohol, other than a few sips	18.3	1,364	2.62	(13.3 - 24.7)	6.3
QN27: Percentage of students who had their first drink of alcohol other than a few sips before age 11 years	6.8	1,355	0.92	(5.0 - 9.1)	1.8
QN28: Percentage of students who ever used marijuana	6.0	1,394	1.33	(3.7 - 9.7)	4.3
QN29: Percentage of students who tried marijuana for the first time before age 11 years	2.9	1,402	1.13	(1.2 - 6.7)	6.4
QN30: Percentage of students who ever used any form of cocaine, including powder, crack, or freebase	2.7	1,461	0.36	(2.0 - 3.6)	0.7
QN31: Percentage of students who ever sniffed glue, breathed the contents of spray cans, or inhaled any paints or sprays to get high	6.3	1,460	0.88	(4.7 - 8.5)	1.9
QN32: Percentage of students who ever took steroid pills or shots without a doctor's prescription	1.3	1,457	0.37	(0.7 - 2.4)	1.6

*The average design effect across all items for this survey is 4.1.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Northern) Middle School Survey Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN33: Percentage of students who ever took a prescription drug (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax) without a doctor's prescription	2.5	1,447	0.43	(1.7 - 3.6)	1.1
QN38: Percentage of students who described themselves as slightly or very overweight	24.3	1,458	1.64	(20.9 - 28.1)	2.1
QN39: Percentage of students who were trying to lose weight	39.3	1,452	2.96	(33.1 - 45.9)	5.3
QN40: Percentage of students who had ever gone without eating for 24 hours or more to lose weight or to keep from gaining weight	15.1	1,450	2.09	(11.1 - 20.2)	5.0
QN41: Percentage of students who had ever taken diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight	2.8	1,458	0.68	(1.6 - 4.7)	2.5
QN42: Percentage of students who had ever vomited or taken laxatives to lose weight or to keep from gaining weight	4.1	1,447	0.66	(2.9 - 5.8)	1.6
QN43: Percentage of students who ate breakfast on all of the past seven days	56.0	1,447	3.67	(47.9 - 63.8)	7.9
QN44: Percentage of students who were physically active for a total of at least 60 minutes per day on five or more of the past seven days	51.9	1,435	2.80	(45.8 - 58.0)	4.5

*The average design effect across all items for this survey is 4.1.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Northern) Middle School Survey Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN45: Percentage of students who watched three or more hours per day of TV on an average school day	25.5	1,424	1.96	(21.5 - 30.0)	2.9
QN46: Percentage of students who played video or computer games or used a computer for something that was not school work three or more hours per day on an average school day	32.8	1,440	2.76	(27.1 - 39.1)	5.0
QN47: Percentage of students who attended physical education (PE) classes one or more days in an average week when they were in school	98.1	1,422	0.54	(96.5 - 99.0)	2.2
QN48: Percentage of students who played on one or more sports teams during the past 12 months	68.2	1,407	3.05	(61.2 - 74.5)	6.1
QN49: Percentage of students who had ever been taught in school about AIDS or HIV infection	42.8	1,421	5.41	(31.6 - 54.7)	17.0
QN50: Percentage of students who had ever been told by a doctor or nurse that they had asthma	18.4	1,417	1.40	(15.5 - 21.6)	1.9
QN51: Percentage of students who smoked Malboro or Newport brand of cigarettes during the past 30 days	0.7	1,454	0.30	(0.3 - 1.8)	1.8
QN52: Percentage of students who usually smoked cigarillos or small cigars during the past 30 days	0.6	1,460	0.27	(0.2 - 1.6)	2.0

*The average design effect across all items for this survey is 4.1.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

**Virginia (Northern) Middle School Survey
Sample Statistics Report**

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN53: Percentage of students who most recently tried smoking tobacco from a hookah or waterpipe, dissolvable tobacco products, or electronic cigarettes	2.0	1,445	0.57	(1.1 - 3.7)	2.4
QN54: Percentage of students who mostly used flavored little cigars or cigarettes or electronic cigarettes during the past 30 days	0.7	1,440	0.21	(0.3 - 1.3)	1.0
QN55: Percentage of students who most of the time or always saw ads or promotions for cigarettes and other tobacco products when they went to a convenience store, supermarket, or gas station	41.1	1,452	3.01	(34.7 - 47.8)	5.4
QN56: Percentage of students who describe the rule inside the home where they live as smoking is allowed at some times or in some places or is always allowed inside their home	10.6	1,446	2.09	(6.8 - 16.1)	6.7
QN57: Percentage of students who describe the rule about smoking in the vehicle they drive or ride in the most as sometimes or always allowed inside the vehicle	14.2	1,440	2.73	(9.2 - 21.3)	8.8
QN58: Percentage of students who drank 100% fruit juices one or more times during the past seven days	79.1	1,453	1.32	(76.0 - 81.8)	1.5
QN59: Percentage of students who ate fruit one or more times during the past seven days	91.9	1,449	1.13	(89.0 - 94.0)	2.5

*The average design effect across all items for this survey is 4.1.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

**Virginia (Northern) Middle School Survey
Sample Statistics Report**

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN60: Percentage of students who ate green salad one or more times during the past seven days	57.0	1,447	2.79	(50.8 - 62.9)	4.6
QN61: Percentage of students who ate potatoes one or more times during the past seven days	58.9	1,452	2.09	(54.3 - 63.4)	2.6
QN62: Percentage of students who ate carrots one or more times during the past seven days	56.8	1,447	2.54	(51.2 - 62.2)	3.8
QN63: Percentage of students who ate other vegetables one or more times during the past seven days	85.0	1,436	1.48	(81.5 - 88.0)	2.5
QN64: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day during the past seven days	13.7	1,449	2.78	(8.7 - 20.9)	9.5
QN65: Percentage of students who drank a can, bottle, or glass of a sugar-sweetened beverage such as lemonade, sweetened tea or coffee drinks, flavored milk, Snapple, or Sunny Delight one or more times per day during the past seven days	12.6	1,448	1.42	(9.8 - 16.0)	2.6
QN66: Percentage of students who drank a can, bottle, or glass of an energy drink, such as Red Bull or Jolt, one or more times per day during the past seven days	3.2	1,439	0.83	(1.8 - 5.6)	3.1

*The average design effect across all items for this survey is 4.1.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

**Virginia (Northern) Middle School Survey
Sample Statistics Report**

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN67: Percentage of students who drank a can, bottle, or glass of a sports drink such as Gatorade or PowerAde one or more times per day during the past seven days	11.0	1,442	1.79	(7.7 - 15.5)	4.7
QN68: Percentage of students who drank a bottle or glass of plain water four or more times per day during the past seven days	34.9	1,441	2.31	(30.0 - 40.1)	3.4
QN69: Percentage of students who drank three or more glasses per day of milk during the past seven days	13.9	1,415	1.13	(11.6 - 16.6)	1.5
QN70: Percentage of students who think young people who smoke definitely or probably have more fiends	26.3	1,430	4.01	(18.5 - 35.9)	11.8
QN71: Percentage of students who say smoking from a hookah or waterpipe is very or somewhat dangerous	91.2	1,410	2.25	(84.9 - 95.1)	8.9
QN72: Percentage of students who say smoking cigars, cigarillos, or little cigars is very or somewhat dangerous	97.2	1,422	0.65	(95.4 - 98.4)	2.2
QN73: Percentage of students who say smoking flavored cigarettes is very or somewhat dangerous to their health	94.5	1,413	1.01	(91.8 - 96.3)	2.8

*The average design effect across all items for this survey is 4.1.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.