

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Southwestern) High School Survey Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN8: Among students who rode a bicycle during the past 12 months, the percentage who never or rarely wore a bicycle helmet	84.6	748	2.33	(78.9 - 89.0)	3.1
QN9: Percentage of students who never or rarely wore a seat belt when riding in a car driven by someone else	9.9	1,203	1.17	(7.7 - 12.8)	1.8
QN10: Percentage of students who rode one or more times during the past 30 days in a car or other vehicle driven by someone who had been drinking alcohol	19.8	1,200	2.07	(15.7 - 24.7)	3.2
QN11: Among students who drove a car or other vehicle during the past 30 days, the percentage who drove when they had been drinking alcohol one or more times during the past 30 days	9.2	678	2.16	(5.4 - 15.1)	3.8
QN12: Among students who drove a car or other vehicle during the past 30 days, the percentage who texted or e-mailed while driving on one or more of the past 30 days	33.5	690	3.99	(25.4 - 42.7)	4.9
QN13: Percentage of students who carried a weapon such as a gun, knife, or club on one or more of the past 30 days	21.9	1,172	1.59	(18.6 - 25.6)	1.7
QN14: Percentage of students who carried a gun on one or more of the past 30 days	10.8	1,184	1.13	(8.6 - 13.5)	1.6

*The average design effect across all items for this survey is 2.3.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Southwestern) High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN16: Percentage of students who did not go to school on one or more of the past 30 days because they felt they would be unsafe at school or on their way to or from school	6.7	1,208	0.95	(4.9 - 9.1)	1.7
QN17: Percentage of students who had been threatened or injured with a weapon such as a gun, knife, or club on school property one or more times during the past 12 months	6.5	1,198	0.63	(5.3 - 8.0)	0.8
QN18: Percentage of students who were in a physical fight one or more times during the past 12 months	21.4	1,195	1.56	(18.2 - 25.0)	1.7
QN19: Percentage of students who were in a physical fight one or more times during the past 12 months in which they were injured and had to be treated by a doctor or nurse	2.6	1,192	0.53	(1.6 - 4.0)	1.4
QN22: Among students who dated or went out with someone during the past 12 months, the percentage who had been physically hurt on purpose by someone they were dating or going out with one or more times during the past 12 months	12.0	812	1.94	(8.4 - 16.9)	2.9
QN24: Percentage of students who had ever been bullied on school property during the past 12 months	24.6	1,202	2.66	(19.3 - 30.9)	4.6
QN25: Percentage of students who had ever been electronically bullied during the past 12 months	17.3	1,201	1.64	(14.0 - 21.1)	2.3

*The average design effect across all items for this survey is 2.3.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Southwestern) High School Survey Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN26: Percentage of students who felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	27.1	1,199	1.97	(23.0 - 31.6)	2.4
QN27: Percentage of students who seriously considered attempting suicide during the past 12 months	15.3	1,184	1.11	(13.0 - 17.8)	1.1
QN28: Percentage of students who made a plan about how they would attempt suicide during the past 12 months	13.3	1,199	1.21	(10.9 - 16.2)	1.5
QN29: Percentage of students who actually attempted suicide one or more times during the past 12 months	8.4	1,066	0.73	(6.9 - 10.1)	0.7
QN30: Percentage of students who made a suicide attempt during the past 12 months that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse	3.2	1,063	0.47	(2.3 - 4.4)	0.8
QN31: Percentage of students who ever tried cigarette smoking, even one or two puffs	42.9	1,158	2.70	(37.1 - 48.8)	3.4
QN32: Percentage of students who smoked a whole cigarette for the first time before age 13 years	10.9	1,140	1.35	(8.3 - 14.2)	2.1
QN33: Percentage of students who smoked cigarettes on one or more of the past 30 days	15.7	1,144	1.93	(12.0 - 20.4)	3.2

*The average design effect across all items for this survey is 2.3.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Southwestern) High School Survey Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN34: Among students who reported current cigarette use, the percentage who smoked more than 10 cigarettes per day on the days they smoked during the past 30 days	14.9	169	4.23	(7.8 - 26.6)	2.4
QN37: Percentage of students who ever smoked cigarettes daily, that is, at least one cigarette every day for 30 days	10.2	1,160	1.24	(7.8 - 13.2)	1.9
QN38: Among students who reported current cigarette use, the percentage who ever tried to quit smoking cigarettes during the past 12 months	50.2	167	3.55	(42.5 - 57.9)	0.8
QN39: Percentage of students who used chewing tobacco, snuff, or dip on one or more of the past 30 days	15.1	1,202	1.10	(12.8 - 17.6)	1.1
QN40: Percentage of students who smoked cigars, cigarillos, or little cigars on one or more of the past 30 days	14.4	1,205	1.87	(10.8 - 19.0)	3.4
QN41: Percentage of students who had at least one drink of alcohol on one or more days during their life	58.4	1,159	3.00	(51.7 - 64.7)	4.3
QN42: Percentage of students who had their first drink of alcohol other than a few sips before age 13 years	19.8	1,187	1.52	(16.8 - 23.4)	1.7
QN43: Percentage of students who had at least one drink of alcohol on one or more of the past 30 days	31.2	1,088	2.83	(25.4 - 37.7)	4.1

*The average design effect across all items for this survey is 2.3.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Southwestern) High School Survey Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN44: Percentage of students who had five or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days	19.9	1,159	2.01	(15.8 - 24.6)	2.9
QN45: Percentage of students who had ten or more drinks of alcohol in a row, that is, within a couple of hours, during the past 30 days	6.7	1,124	1.80	(3.7 - 11.8)	5.8
QN46: Among students who reported current alcohol use, the percentage who usually got the alcohol they drank from someone who gave it to them during the past 30 days	41.0	319	3.68	(33.3 - 49.2)	1.8
QN47: Percentage of students who used marijuana one or more times during their life	31.6	1,173	2.70	(26.0 - 37.8)	4.0
QN48: Percentage of students who tried marijuana for the first time before age 13 years	8.6	1,175	1.02	(6.7 - 11.1)	1.5
QN49: Percentage of students who used marijuana one or more times during the past 30 days	18.9	1,180	1.70	(15.5 - 22.9)	2.2
QN50: Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during their life	7.5	1,201	0.84	(5.9 - 9.5)	1.2

*The average design effect across all items for this survey is 2.3.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Southwestern) High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN51: Percentage of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during their life	8.8	1,201	0.73	(7.4 - 10.6)	0.8
QN52: Percentage of students who used heroin one or more times during their life	4.4	1,192	0.82	(2.9 - 6.6)	1.9
QN53: Percentage of students who used methamphetamines one or more times during their life	4.4	1,200	0.78	(3.0 - 6.5)	1.7
QN54: Percentage of students who used ecstasy one or more times during their life	7.1	1,200	0.79	(5.6 - 9.1)	1.1
QN55: Percentage of students who took steroid pills or shots without a doctor's prescription one or more times during their life	3.5	1,194	0.64	(2.4 - 5.2)	1.4
QN56: Percentage of students who have taken a prescription drug (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax) without a doctor's prescription one or more times during their life	15.9	1,197	1.40	(13.1 - 19.2)	1.7
QN57: Percentage of students who used a needle to inject any illegal drug into their body one or more times during their life	3.7	1,192	0.63	(2.5 - 5.4)	1.3
QN66: Percentage of students who described themselves as slightly or very overweight	33.4	1,189	1.76	(29.6 - 37.3)	1.7

*The average design effect across all items for this survey is 2.3.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

**Virginia (Southwestern) High School Survey
Sample Statistics Report**

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN67: Percentage of students who were trying to lose weight	43.6	1,194	1.86	(39.6 - 47.7)	1.7
QN68: Percentage of students who went without eating for 24 hours or more to lose weight or to keep from gaining weight during the past 30 days	12.8	1,176	0.67	(11.4 - 14.3)	0.5
QN69: Percentage of students who took any diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight during the past 30 days	7.0	1,182	1.07	(5.0 - 9.7)	2.1
QN70: Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight during the past 30 days	6.5	1,179	0.59	(5.3 - 7.9)	0.7
QN71: Percentage of students who drank 100% fruit juices one or more times during the past seven days	70.2	1,196	0.66	(68.8 - 71.6)	0.2
QN72: Percentage of students who ate fruit one or more times during the past seven days	82.9	1,193	1.53	(79.3 - 85.9)	2.0
QN73: Percentage of students who ate green salad one or more times during the past seven days	55.2	1,196	1.93	(50.9 - 59.3)	1.8
QN74: Percentage of students who ate potatoes one or more times during the past seven days	69.2	1,195	1.26	(66.4 - 71.8)	0.9
QN75: Percentage of students who ate carrots one or more times during the past seven days	42.3	1,190	1.97	(38.1 - 46.6)	1.9

*The average design effect across all items for this survey is 2.3.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Southwestern) High School Survey Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN76: Percentage of students who ate other vegetables one or more times during the past seven days	79.0	1,193	1.37	(75.9 - 81.9)	1.3
QN77: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day during the past seven days	34.3	1,192	2.35	(29.4 - 39.6)	2.9
QN78: Percentage of students who drank three or more glasses per day of milk during the past seven days	9.0	1,184	0.92	(7.2 - 11.2)	1.2
QN79: Percentage of students who ate breakfast on all of the past seven days	38.9	1,182	2.08	(34.5 - 43.5)	2.1
QN80: Percentage of students who were physically active for a total of at least 60 minutes per day on five or more of the past seven days	47.3	1,171	2.21	(42.6 - 52.1)	2.3
QN81: Percentage of students who watched three or more hours per day of TV on an average school day	29.3	1,180	2.42	(24.4 - 34.9)	3.3
QN82: Percentage of students who played video or computer games or used a computer for something that was not school work three or more hours per day on an average school day	40.2	1,180	2.27	(35.4 - 45.3)	2.5
QN83: Percentage of students who attended physical education (PE) classes on one or more days in an average week when they were in school	39.5	1,178	3.69	(31.8 - 47.7)	6.7

*The average design effect across all items for this survey is 2.3.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Southwestern) High School Survey Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN84: Percentage of students who played on one or more sports teams during the past 12 months	56.3	1,170	2.63	(50.5 - 61.9)	3.3
QN85: Percentage of students who had ever been taught in school about AIDS or HIV infection	81.0	1,169	3.12	(73.3 - 86.9)	7.4
QN86: Percentage of students who had ever been told by a doctor or nurse that they had asthma	24.3	1,169	2.38	(19.5 - 29.8)	3.6
QN87: Percentage of students who have ever been bullied away from school property during the past 12 months	16.7	1,191	1.46	(13.7 - 20.1)	1.8
QN88: Percentage of students who have ever bullied someone else on school property during the past 12 months	12.7	1,202	1.41	(9.9 - 16.1)	2.2
QN89: Percentage of students who have ever bullied someone else away from school property during the past 12 months	10.4	1,202	0.90	(8.6 - 12.5)	1.1
QN90: Percentage of students who have ever been the victim of teasing or name calling during the past 12 months because of their weight, size, or physical appearance	29.8	1,192	1.64	(26.4 - 33.5)	1.5
QN91: Percentage of students who have ever been the victim of teasing or name calling during the past 12 months because of their gender	7.8	1,200	0.85	(6.1 - 9.8)	1.2

*The average design effect across all items for this survey is 2.3.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Southwestern) High School Survey Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN92: Percentage of students who smoked Marlboro or Newport brand of cigarettes during the past 30 days	12.2	1,198	2.28	(8.1 - 18.1)	5.8
QN93: Percentage of students who usually smoked cigarillos or small cigars during the past 30 days	8.5	1,200	1.83	(5.3 - 13.4)	5.1
QN94: Percentage of students who most recently tried smoking tobacco from a hookah or waterpipe, dissolvable tobacco products, or electronic cigarettes	7.8	1,185	1.05	(5.8 - 10.4)	1.8
QN95: Percentage of students who mostly used flavored little cigars or cigarettes or electronic cigarettes during the past 30 days	8.5	1,177	1.67	(5.5 - 12.9)	4.2
QN96: Percentage of students who most of the time or always saw ads or promotions for cigarettes and other tobacco products when they went to a convenience store, supermarket, or gas station	70.2	1,201	2.00	(65.6 - 74.3)	2.3
QN97: Percentage of students who describe the rule about smoking inside the home where they live as smoking is allowed at some times or in some places or smoking is always allowed inside their home	26.3	1,193	2.01	(22.2 - 30.9)	2.5
QN98: Percentage of students who describe the rule about smoking in the vehicle they drive or ride in the most as smoking is sometimes allowed inside the vehicle or smoking is always allowed inside the vehicle	37.9	1,184	2.47	(32.7 - 43.4)	3.1

*The average design effect across all items for this survey is 2.3.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Southwestern) High School Survey Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN99: Percentage of students who drank a can, bottle, or glass of a sugar-sweetened beverage such as lemonade, sweetened tea or coffee drinks, flavored milk, Snapple, or Sunny Delight one or more times per day during the past seven days	29.1	1,181	2.09	(24.7 - 33.8)	2.5
QN100: Percentage of students who drank a can, bottle, or glass of an energy drink, such as Red Bull or Jolt, one or more times per day during the past seven days	5.5	1,195	0.63	(4.3 - 7.0)	0.9
QN101: Percentage of students who drank a can, bottle, or glass of a sports drink such as Gatorade or PowerAde one or more times per day during the past seven days	15.0	1,196	1.39	(12.2 - 18.3)	1.8
QN102: Percentage of students who drank a bottle or glass of plain water four or more times per day during the past seven days	25.3	1,183	1.45	(22.3 - 28.6)	1.3
QN103: Percentage of students who walk or ride their bike to school on five days in an average week when they are in school and when the weather allows	3.7	1,162	0.35	(3.0 - 4.6)	0.4
QN104: Percentage of students who think young people who smoke definitely or probably have more friends	22.8	1,154	1.49	(19.7 - 26.2)	1.5
QN105: Percentage of students who say smoking from a hookah or waterpipe is very or somewhat dangerous to their health	89.1	1,156	1.27	(86.0 - 91.6)	1.9

*The average design effect across all items for this survey is 2.3.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Virginia (Southwestern) High School Survey Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN106: Percentage of students who say smoking cigars, cigarillos, or little cigars is very or somewhat dangerous to their health	95.1	1,172	0.79	(93.1 - 96.6)	1.6
QN107: Percentage of students who say smoking flavored cigarettes is very or somewhat dangerous to their health	95.6	1,165	0.76	(93.6 - 97.0)	1.6
QN108: Percentage of students who most of the time or always went hungry during the past 30 days because there was not enough food in their home	3.7	1,174	0.73	(2.4 - 5.7)	1.8
QN109: Percentage of students who responded that there is at least one teacher or other adult in this school that they can talk to if they have a problem	74.5	1,166	1.63	(70.8 - 77.9)	1.6
QN110: Percentage of students who most of the time or always feel safe and secure at school	74.0	1,175	1.71	(70.1 - 77.6)	1.8
QN111: Percentage of students who strongly agree or agree that their school has clear rules and consequences for behavior	62.8	1,171	2.00	(58.4 - 67.1)	2.0
QN112: Percentage of students who disagree or strongly disagree that harassment and bullying by other students is a problem at their school	21.9	1,176	2.95	(16.2 - 29.0)	6.0
QN113: Percentage of students who disagree or strongly disagree that violence is a problem at their school	36.9	1,169	3.02	(30.6 - 43.7)	4.6

*The average design effect across all items for this survey is 2.3.

Note: Design effect is calculated differently for the 2013 Sample Statistics Report than it has been in the past. For 2013, the design effect for each question is calculated by dividing the variance for each question by the variance that would have resulted from a simple random sample. In previous years, design effect was calculated by dividing the standard error for each question by the standard error that would have resulted from a simple random sample. Consequently, design effects for 2013 will be larger than in the past even though no changes have been made to the standard sample design.