

Smoking and Pregnancy

Why are we concerned about women smoking while they are pregnant?

Smoking during pregnancy is a common cause of pregnancy complications and of illness and death among infants. Smoking can lead to problems for a woman trying to get pregnant or who is already pregnant, and for her baby before and after birth.

What are the effects of smoking on pregnant women?

- Increased risk for infections and chronic illnesses such as emphysema, heart disease, and lung cancer
- Placenta separates from the womb too early, causing bleeding
- Placenta covers the cervix, causing complications
- Water breaks too early
- Miscarriage

What are the effects of smoking on babies?

- Damage unborn baby's growing brain and lungs
- Baby born too small or too early
- Baby born with cleft lip or cleft palate
- Stillbirth or infant death
- Sudden Infant Death Syndrome (SIDS)
- Possible cognitive effects associated with learning disabilities and conduct disorders



Where can I get help for quitting smoking?

Quitting smoking can be hard, but it is one of the best ways a woman can protect herself and her baby's health. For support in quitting, including free quit coaching, a free quit plan, free education materials, and referrals to local resources, call **1-800-Quit-Now** (1-800-784-8669); TTY: 1-800-332-8615. Services are available for free in English and Spanish.

For more information: www.quitnow.net/virginia or www.cdc.gov/reproductivehealth/maternalinfanthealth/tobaccousepregnancy



1/16

Health  Whys

A public health message from
www.tjhd.org

Follow us on
Facebook



VDH VIRGINIA
DEPARTMENT
OF HEALTH
Protecting You and Your Environment

You can find this and other timely public health tips and events on the Thomas Jefferson Health District (TJHD) Facebook page. Check us out and "like us!"