

Table 5. Reported Total Early Syphilis (TES)* Cases^ by Quarter, Year of Report and Selected Demographics


<u>Sex</u>	Quarter	2007					2008					2009		
		1	2	3	4	Total	1	2	3	4	Total	1	2	Total
Female		15	14	9	20	58	17	14	12	14	57	17	17	34
Male		80	73	102	94	349	108	120	110	104	442	87	125	212
Total		95	87	111	114	407	125	134	122	118	499	104	142	246
<u>Race/Ethnicity</u>														
Asian/Pacific Islander		1	1	2	0	4	3	1	3	1	8	0	0	0
Black		46	45	61	73	225	77	89	83	87	336	76	95	171
Hispanic		6	8	6	8	28	6	5	5	8	24	8	10	18
White		42	33	42	33	150	39	37	31	18	125	19	34	53
Other/Not Reported		0	0	0	0	0	0	2	0	4	6	1	3	4
Total		95	87	111	114	407	125	134	122	118	499	104	142	246
<u>Age at diagnosis</u>														
10 - 14		0	0	0	1	1	0	0	0	0	0	0	1	1
15 - 19		6	9	9	8	32	9	10	10	13	42	13	12	25
20 - 29		33	30	26	41	130	49	50	50	53	202	46	75	121
30 - 39		22	21	36	31	110	26	30	26	25	107	22	19	41
40 - 49		24	22	26	25	97	25	32	24	16	97	16	27	43
50+		10	5	14	8	37	16	12	12	11	51	7	8	15
Total		95	87	111	114	407	125	134	122	118	499	104	142	246

* Total Early Syphilis includes: Primary, Secondary and Early Latent Syphilis

^ See definitions under [Technical Notes](#)

Gender Proportions


Cumulative for Qtrs in Table



Female 12.9%
Male 87.1%
Total: 100.0%

Race Proportions

Cumulative for Qtrs in Table



Asian/Pacific Islander 1.04%
Black 63.54%
Hispanic 6.08%
Other/Not Reported 0.87%
White 28.47%
Total: 100.00%