

Table 5a. Reported Total Early Syphilis (TES)* Cases^ by Race and Select Demographics, Year to Date^

	Asian/Pacific Islander		Black		Hispanic		White		Other/Not Reported		Total	
Sex												
Female	0	0%	45	13%	2	8%	10	8%	0	0%	57	11%
Male	8	100%	292	86%	22	91%	115	92%	6	100%	443	88%
All	8	100%	337	100%	24	100%	125	100%	6	100%	500	100%
Age at diagnosis												
15 - 19	0	0%	39	11%	2	8%	1	0%	0	0%	42	8%
20 - 29	2	25%	159	47%	7	29%	31	24%	3	50%	202	40%
30 - 39	4	50%	70	20%	10	41%	22	17%	1	16%	107	21%
40 - 49	1	12%	52	15%	3	12%	41	32%	1	16%	98	19%
50+	1	12%	17	5%	2	8%	30	24%	1	16%	51	10%
All	8	100%	337	100%	24	100%	125	100%	6	100%	500	100%
Region												
Central	1	12%	127	37%	3	12%	43	34%	2	33%	176	35%
Eastern	2	25%	141	41%	4	16%	25	20%	4	66%	176	35%
Northern	5	62%	39	11%	17	70%	40	32%	0	0%	101	20%
Northwest	0	0%	6	1%	0	0%	10	8%	0	0%	16	3%
Southwest	0	0%	24	7%	0	0%	7	5%	0	0%	31	6%
All	8	100%	337	100%	24	100%	125	100%	6	100%	500	100%

* Total Early Syphilis includes: Primary, Secondary and Early Latent Syphilis

^ See definitions under [Technical Notes](#)

Reported Cases of TES by Region, Year to Date

