

Table 5a. Reported Total Early Syphilis (TES)* Cases^ by Race and Selected Demographics, Year to Date^

Sex	Black		Hispanic		White		Other/Not Reported		Total	
	Count	%	Count	%	Count	%	Count	%	Count	%
Female	21	12%	3	17%	9	17%	1	25%	34	14%
Male	150	88%	15	83%	44	83%	3	75%	212	86%
All	171	100%	18	100%	53	100%	4	100%	246	100%

Age at diagnosis

Age Group	Black		Hispanic		White		Other/Not Reported		Total	
	Count	%	Count	%	Count	%	Count	%	Count	%
10 - 14	1	1%	0	0%	0	0%	0	0%	1	0%
15 - 19	21	12%	1	6%	3	6%	0	0%	25	10%
20 - 29	87	51%	9	50%	23	43%	2	50%	121	49%
30 - 39	28	16%	4	22%	7	13%	2	50%	41	17%
40 - 49	24	14%	3	17%	16	30%	0	0%	43	17%
50+	10	6%	1	6%	4	8%	0	0%	15	6%
All	171	100%	18	100%	53	100%	4	100%	246	100%

Region


Region	Black		Hispanic		White		Other/Not Reported		Total	
	Count	%	Count	%	Count	%	Count	%	Count	%
Central	54	32%	6	33%	9	17%	0	0%	69	28%
Eastern	71	42%	2	11%	18	34%	2	50%	93	38%
Northern	17	10%	9	50%	16	30%	1	25%	43	17%
Northwest	4	2%	0	0%	6	11%	0	0%	10	4%
Southwest	25	15%	1	6%	4	8%	1	25%	31	13%
All	171	100%	18	100%	53	100%	4	100%	246	100%

* Total Early Syphilis includes: Primary, Secondary and Early Latent Syphilis

^ See definitions under [Technical Notes](#)

Figure 5a. Reported Total Early Syphilis (TES)* Cases^ by Race and Selected Demographics, Year to Date^

Race/Ethnicity and Gender



By Race/Ethnicity and Region

