

Arthritis by Demographics (Race/Ethnicity, Education, Income, Age, and Gender), Virginia, 2011

		Have you ever been told that you have some form of arthritis, rheumatoid arthritis, gout, lupus, or fibromyalgia?							
		Yes				No			
		Sample Size	Weighted Counts	Weighted Percent (%)	CI	Sample Size	Weighted Counts	Weighted Percent (%)	CI
*Virginia	*State Total	2238	1576468	25.7	24.3 -27.1	4327	4550214	74.3	72.9 - 75.7
Race/Ethnicity	Black/Non-Hispanic	330	250134	22.5	19.3 -25.8	628	860521	77.5	74.2 - 80.7
	Hispanic	34	33353	8.1	5.0 - 11.3	219	376125	91.9	88.7 - 95.0
	Other	85	88257	20.5	14.8 -26.2	232	342097	79.5	73.8 - 85.2
	White/Non-Hispanic	1750	1176289	28.7	27.0 -30.4	3193	2919674	71.3	69.6 - 73.0
Education	< H.S.	321	337712	40.1	35.0 -45.1	278	505147	59.9	54.9 - 65.0
	H.S. or G.E.D.	696	432078	26.4	23.8 -28.9	1101	1205980	73.6	71.1 - 76.2
	Some College	567	443681	25.6	22.7 -28.5	1016	1288895	74.4	71.5 - 77.3
	College Graduate	642	359128	19.0	17.0 -21.0	1904	1528244	81.0	79.0 - 83.0
Income	\$15,000 or less	299	210572	46.1	39.9 -52.4	219	246019	53.9	47.6 - 60.1
	\$15,000 to less than \$25,000	413	244407	27.6	24.2 -30.9	566	642663	72.4	69.1 - 75.8
	\$25,000 to less than \$35,000	226	157984	28.9	24.2 -33.6	370	388235	71.1	66.4 - 75.8
	\$35,000 to less than \$50,000	280	190266	27.0	22.9 -31.2	513	513798	73.0	68.8 - 77.1
	\$50,000 or more	660	535132	19.9	17.9 -21.9	2062	2150245	80.1	78.1 - 82.1
Age	Age 18 to 24	---	---	---	---	241	750019	95.1	92.4 - 97.7
	Age 25 to 34	54	71713	6.8	4.5 - 9.1	699	982334	93.2	90.9 - 95.5
	Age 35 to 44	137	178445	16.0	13.0 -19.1	749	934921	84.0	80.9 - 87.0

	Age 45 to 54	360	330499	27.7	24.3 -31.2	889	861452	72.3	68.8 -
	Age 55 to 64	657	407230	42.6	39.2 -46.0	880	548808	57.4	54.0 -
	Age 65 or older	1016	549580	53.8	50.5 -57.0	869	472680	46.2	43.0 -
Gender	Female	1497	959269	30.5	28.5 -32.5	2436	2184817	69.5	67.5 -
	Male	741	617199	20.7	18.8 -22.6	1891	2365397	79.3	77.4 -

1. Data source: Virginia Department of Health, Division of Policy and Evaluation, Behavioral Risk Factor Surveillance Survey, 2011. Weighted counts and weighted percents are weighted to population characteristics.
2. "-" replaces estimates when the unweighted sample size for the denominator was < 20 or the CI half width was > 10 for any cell.
3. Responses of don't know/not sure, refused, or missing were removed from the numerator and denominator in all estimates.
4. \*\*\* Use caution in interpreting sample sizes less than 50.
5. CI= 95% confidence interval.
6. \*\*\* Arthritis diagnoses include: rheumatism, polymyalgia rheumatica, osteoarthritis (not osteoporosis), tendonitis, bursitis, bunion, tennis elbow, carpal tunnel syndrome, tarsal tunnel syndrome, joint infection, Reiter's syndrome, ankylosing spondylitis, spondylosis, rotator cuff syndrome, connective tissue disease, scleroderma, polymyositis, Raynaud's syndrome, vasculitis (giant cell arteritis, Henoch-Schonlein purpura, Wegener's granulomatosis, polyarteritis nodosa)