



EPIDEMIOLOGY BULLETIN

James B. Kenley, M.D., Commissioner
Grayson B. Miller, Jr., M.D., Epidemiologist

Editor: Carl W. Armstrong, M.D.

November, 1983

Volume 83, Number 11

Yellow Fever Vaccine

Recommendations of the Immunization Practices Advisory Committee of the U.S. Public Health Service

Introduction

Yellow fever presently occurs only in Africa and South America. Two forms of yellow fever—urban and jungle—are epidemiologically distinguishable. Clinically and etiologically, they are identical^{1,2}.

Urban yellow fever is an epidemic viral disease of humans transmitted from infected to susceptible persons by a vector, the *Aedes aegypti* mosquito. In areas where *Ae. aegypti* has been eliminated or suppressed, urban yellow fever has disappeared; eradication of *Ae. aegypti* in a number of countries, notably Panama, Brazil, Ecuador, Peru, Bolivia, Paraguay, Uruguay, and Argentina, achieved in the early 1900s, led to the disappearance of urban yellow fever. The last *Ae. aegypti*-borne yellow fever epidemic occurred in Trinidad in 1954. However, periodic reinfestations of some countries have occurred in recent years, and other countries remain infested, including areas of Venezuela, Colombia, and Guiana, which border on the enzootic zone for jungle yellow fever. In West Africa, *Ae. aegypti*-transmitted epidemics continue to occur at frequent intervals and involve human populations in both towns and rural villages³.

Jungle yellow fever is an enzootic viral disease transmitted among non-human primate hosts by a variety of mosquito vectors. It is currently observed only in forest-savannah zones of tropical Africa and in forested areas of South America, but occasionally extends into parts of Central America and the island of Trinidad. In South

America, approximately 200-400 cases are recognized annually, mainly among persons with occupational exposures in forested areas; the disease is, however, believed to be greatly underreported. In Africa, epidemics involving forest mosquito vectors affect tens of thousands of persons at intervals of a few years, but few cases are officially reported. The disease may sometimes not be detected in an area for some years and then reappear. Delineation of affected areas depends on surveillance of animal reservoirs and vectors, accurate diagnosis, and prompt reporting of all cases. The jungle yellow fever cycle may be ac-


tive but unrecognized in forested areas of countries within the yellow fever endemic zone (Figure 1).

Urban yellow fever can be prevented by eradicating *Ae. aegypti* mosquitoes or by suppressing their numbers to the point that they no longer perpetuate infection. At the present time, jungle yellow fever can most effectively be prevented in humans by immunization.

Yellow Fever Vaccine

Yellow fever vaccine* is a live, attenuated virus preparation made from the 17D yellow fever virus strain⁴. The

*Official name: Yellow Fever Vaccine.
Continued on page 2



Continued from page 1

17D vaccine has proven to be extremely safe and effective⁵. The 17D strain is grown in chick embryo inoculated with a seed virus of a fixed-passage level. The vaccine is freeze-dried supernate of centrifuged embryo homogenate, packaged in one-dose and five-dose vials for domestic use.

Vaccine should be stored at temperatures between 5 C (41 F) and -30 C (-22 F)—preferably frozen, below 0 C (32 F)—until it is reconstituted by the addition of diluent sterile, physiologic saline supplied by the manufacturer. Multiple dose vials of reconstituted vaccine should be held at 5 C-10 C (41 F-50 F); unused vaccine should be discarded within 1 hour after reconstitution.

VACCINE USAGE

A. Persons living or traveling in endemic areas:

1. Persons 6 months of age or older traveling or living in areas where yellow fever infection exists—currently parts of Africa and South America—should be vaccinated. (These are listed in the "Bi-Weekly Summary of Countries with Areas Infected with Quarantinable Diseases" available in state and local health departments. Information on known or probable infected areas is also available from the World Health Organization [WHO] and Pan American Health Organization offices or the Division of Vector-Borne Viral Diseases, Center for Infectious Diseases, CDC, Fort Collins, Colorado.)

Vaccination is also recommended for travel outside the urban areas of countries in the yellow fever endemic zone (Figure 1). It should be emphasized that the actual areas of yellow fever virus activity far exceed the infected zones officially reported and that, in recent years, fatal cases of yellow fever have occurred in unvaccinated tourists⁶.

2. Infants under 6 months of age and pregnant women should be considered for vaccination if traveling to high-risk areas when travel cannot be postponed and a high level of prevention against mosquito exposures is not feasible.

3. Laboratory personnel who might be exposed to virulent yellow fever virus should also be vaccinated.

B. Vaccination for international travel: For purposes of international travel, yellow fever vaccines pro-

duced by different manufacturers worldwide must be approved by WHO and administered at an approved Yellow Fever Vaccination Center. State and territorial health departments have the authority to designate nonfederal vaccination centers; these can be identified by contacting state or local health departments. Vaccinees should have an International Certificate of Vaccination filled in, signed, and validated with the center's stamp where the vaccine is given.

Vaccination for International travel may be required under circumstances other than those specified herein. Some countries in Africa require evidence of vaccination from all entering travelers. Some countries may waive the requirements for travelers coming from noninfected areas and staying less than 2 weeks. These requirements may change, so all travelers should seek current information from health departments. Travel agencies, international airlines, and/or shipping lines should also have up-to-date information.

Some countries require an individual, even if only in transit, to have a valid International Certificate of Vaccination if he or she has been in countries either known or thought to harbor yellow fever virus. Such requirements may be strictly enforced, particularly for persons traveling from Africa or South America to Asia.

C. Primary immunization: For persons of all ages, a single subcutaneous injection of 0.5 ml of reconstituted vaccine is used.

D. Booster doses: Yellow fever immunity following vaccination with 17D strain virus persists for more than 10 years⁷⁻⁹; the International Health Regulations do not require vaccination more often than every 10 years.

REACTIONS

Reactions to 17D yellow fever vaccine are generally mild. Two percent to 5% of vaccinees have mild headaches, myalgia, low-grade fevers, or other minor symptoms 5-10 days after vaccination. Fewer than 0.2% curtail regular activities. Immediate hypersensitivity reactions, characterized by rash, urticaria, and/or asthma, are extremely uncommon (incidence less than 1/1,000,000) and occur principally in persons with histories of egg allergy. Although more than 34 million doses of vaccines have been distrib-

uted, only two cases of encephalitis temporally associated with vaccinations have been reported in the United States; in one fatal case, 17D virus was isolated from the brain.

PRECAUTIONS AND CONTRAINDICATIONS

A. Age: Infants under 6 months of age are theoretically more susceptible to serious adverse reactions (encephalitis) than older children.

B. Pregnancy: Although specific information is not available concerning adverse effects of yellow fever vaccine on the developing fetus, it is prudent on theoretical grounds to avoid vaccinating pregnant women and to postpone travel to areas where yellow fever is present until after delivery. If international travel requirements constitute the only reason to vaccinate a pregnant woman, rather than an increased risk of infection, efforts should be made to obtain a waiver letter from the traveler's physician (see below). Pregnant women who must travel to areas where the risk of yellow fever is high should be vaccinated. It is believed that un-

References

1. Strode GK, ed. Yellow Fever. New York: McGraw-Hill, 1971.
2. World Health Organization. Report of the WHO Expert Committee on Yellow Fever. WHO Tech Rep: Ser. No. 4791, 1971.
3. Monath TP, Craven RB, Adjuikiewicz J. Yellow fever virus. In: Murray PR, Baron EJ, Tenover FC, Tenover FC, eds. Manual of clinical microbiology. Washington, DC: ASM Press, 1979; epidemiologic aspects with other arbovirus infections. Am J Trop Med Hyg 1979; 20:1-10.
4. Smithburn KC, Durieux C, Koerber J. Yellow fever virus. In: Monographs of the World Health Organization. Geneva, 1954; Monograph Series No. 30.
5. Wissemann CL Jr, Sweet BH. Immunity to yellow fever virus. III. Response of human beings to yellow fever vaccine. Am J Trop Med Hyg 1963; 12:411-5.
6. Rodhain F, Hannoun C, Jousset FX. Yellow fever (jaune à Paris à partir de deux cas humains). J Hyg (Paris) 1972; 72:411-5.
7. Groot H, Ribeiro RB. Neutralizing antibody to yellow fever 17 years after vaccination. J Hyg (Paris) 1972; 72:699-707.
8. Poland JD, Calisher CH, Monath TP. Persistence of neutralizing antibody 30-35 years after vaccination. Bull WHO 1981; 59:895-900.
9. Rosenzweig EC, Babione RW, Wissinger R. Yellow fever virus. IV. Response of human beings to yellow fever vaccine following vaccination with 17D strain. J Hyg (Paris) 1963; 12:230-5.
10. Tauraso NM, Myers MG, Nau EV, et al. Persistence of live smallpox and yellow-fever vaccines. J Hyg (Paris) 1972; 126:363-371.
11. Felsenfeld O, Wolf RH, Gyr K, et al. Yellow fever and yellow fever. Lancet 1973; i:1457-8.
12. Gateff C. Acquisitions récentes de la pathologie exotique. Pathol Exot 1972; 65:784-96.
13. Kaplan JE, Nelson DB, Schonberger L. Yellow fever vaccine. In: Trivalent oral polio and yellow fever vaccine.

der these circumstances, the small theoretical risk for mother and fetus from vaccination is far outweighed by the risk of yellow fever infection.

C. *Altered immune states*: Infection with yellow fever vaccine virus poses a theoretical risk to patients with leukemia, lymphoma, or generalized malignancy or to those whose immunologic responses are suppressed by corticosteroids, alkylating drugs, antimetabolites, or radiation. Short-term (less than 2 weeks) corticosteroid therapy or intra-articular, bursal, or tendon injections with corticosteroids should not be immunosuppressive and constitute no increased hazard to recipients of yellow fever vaccine.

D. *Hypersensitivity*: Live yellow fever vaccine is produced in chick embryos and should not be given to persons clearly hypersensitive to eggs; generally, persons who are able to eat eggs or egg products may receive the vaccine.

If international travel regulations are the only reason to vaccinate a patient hypersensitive to eggs, ef-

orts should be made to obtain a waiver. A physician's letter clearly stating the contraindication to vaccination has been acceptable to some governments. (Ideally, it should be written on letterhead stationery and bear the stamp used by health departments and official immunization centers to validate the international Certificates of Vaccination.) Under these conditions, it is also useful for the traveler to obtain specific and authoritative advice from the country or countries he or she plans to visit. Their embassies or consulates may be contacted. Subsequent waiver of requirements should be documented by appropriate letters.

If vaccination of an individual with a questionable history of egg hypersensitivity is considered essential because of a high risk of exposure, an intradermal test dose may be administered under close medical supervision. Specific directions for skin testing are found in the package insert.

Simultaneous Administration of Other Vaccines

Determination of whether to administer yellow fever vaccine and other immunobiologics simultaneously should be made on the basis of convenience to the traveler in completing the desired immunizations before travel and on information regarding possible interference. The following will help guide these decisions.

Studies have shown that the serologic response to yellow fever vaccine is not inhibited by administration of certain other vaccines concurrently or at various intervals of a few days to 1 month. Measles, smallpox, and yellow fever vaccines have been administered in combination with full efficacy of each of the components; Bacillus Calmette Guérin (BCG) and yellow fever vaccines have been administered simultaneously without interference. Additionally, severity of reactions to vaccination was not amplified by concurrent administration of yellow fever and other live virus vaccines¹⁰. If live virus vaccines are not given concurrently, 4 weeks should be allowed to elapse between sequential vaccinations.

Other studies have indicated that persons given yellow fever and cholera vaccines simultaneously or 1-3 weeks apart showed reduced antibody responses to both vaccines^{11,12}. When feasible, cholera and yellow fever vaccines should be administered at a minimal interval of 3 weeks, un-

Veterinarians To Receive *Bulletin*

We are pleased to be able to distribute the *Bulletin* to licensed veterinarians in Virginia. We hope this action will improve communication between the Division of Epidemiology and veterinarians (and vice versa) given the continuing importance to public health of the zoonoses.

The addition of 787 veterinarians to the *Bulletin's* distribution list brings the readership total to 13,000, including close to 10,000 licensed physicians in Virginia, 1,500 local and state health department personnel, health officials in 52 states and territories, 142 hospital infection control practitioners, and 750 libraries, school infirmaries, employee health services, emergency rooms, federal agencies, and individuals who have requested subscription.

Concerned about hospital hazards? Here is a conference for you.

"Hospitals: An Occupational Health and Safety Challenge"
April 11-12, 1984—Richmond

Sponsored by: Bureau of Occupational Health, Virginia State Health Dept. and The Occupational Safety and Health Resource Center
University of North Carolina at Chapel Hill

Designed for hospital occupational health and safety teams (OHN, MD, IH, management, safety director, risk managers).

Contact: Nancy Jakubec
804/281-9527

less time constraints preclude this. If the vaccines cannot be administered at least 3 weeks apart, they should be given simultaneously. There are no data on possible interference between yellow fever and typhoid, paratyphoid, typhus, hepatitis B, plague, rabies, or Japanese encephalitis vaccines.

A recently completed prospective study of persons given yellow fever vaccine and 5 cc of commercially available immune globulin revealed no alteration of the immunologic response to yellow fever vaccine when compared to controls¹³.

Reprinted from *MMWR* 1984; 32: 679-82, 687-8.

ork: McGraw-Hill, 1951.
Com on Yellow Fever, 3rd Report

A, et al. Yellow fever in the Gambia, 1978-
servations on the occurrence of Orungo
29: 912-28, 1980.

R, et al. Yellow fever vaccination. WHO
1956.

ological studies with group B arthropod-
subjects to revaccination with 17D strain
1962; 11:570-5.

Ravisse P. Isôlement du virus de la fièvre
ains importés. Bull Soc Pathol Exot 1979;

nd haemagglutination-inhibiting antibodies
ation with 17D vaccine. Bull WHO 1962;

, Downs WG, Murphy K. Persistence of
ter immunization with 17D yellow fever

eman CL Jr. Immunological studies with
7. Persistence of yellow fever antibodies
n yellow fever vaccine. Am J Trop Med

t al. Effect of interval between inoculation
cines on antigenicity in man. J Infect Dis

Simultaneous vaccination against cholera
8.

atie associations vaccinales. Bull Soc

LB, et al. The effect of immune globulin on
vaccinations. Bull WHO 1984 (in press).

Month: November, 1983

Disease	State					Regions				
	This Month	Last Month	Total to Date		Mean 5 Year To Date	This Month				
			1983	1982		N.W.	N.	S.W.	C.	E.
Measles	0	0	23	14	695	0	0	0	0	0
Mumps	3	2	35	39	103	0	0	1	1	1
Pertussis	0	4	49	28	15	0	0	0	0	0
Rubella	0	1	3	12	102	0	0	0	0	0
Meningitis—Aseptic	50	64	313	247	221	21	2	4	7	16
Other Bacterial	18	17	218	193	162	2	2	3	4	7
Hepatitis A (Infectious)	6	11	118	176	231	1	1	0	2	2
B (Serum)	31	49	494	474	430	2	11	1	6	11
Non-A, Non-B	5	9	75	78	*50	1	1	1	1	1
Salmonellosis	99	185	1,364	1,380	1,200	17	12	15	26	29
Shigellosis	39	32	208	149	374	2	2	0	1	34
Campylobacter Infections	41	66	510	387	*210	10	8	4	6	13
Tuberculosis	39	63	466	570	—	—	—	—	—	—
Syphilis (Primary & Secondary)	35	57	527	579	526	1	6	1	9	18
Gonorrhea	1,676	2,445	19,413	20,001	20,650	—	—	—	—	—
Rocky Mountain Spotted Fever	0	5	61	74	94	0	0	0	0	0
Rabies in Animals	26	44	590	651	171	7	19	0	0	0
Meningococcal Infections	7	9	77	67	73	2	2	0	0	3
Influenza	1	5	902	382	2,306	1	0	0	0	0
Toxic Shock Syndrome	1	0	7	7	*8	1	0	0	0	0
Reyes Syndrome	0	1	6	5	13	0	0	0	0	0
Legionellosis	4	1	24	22	18	2	0	0	1	1
Kawasaki's Disease	1	1	36	14	17	0	0	1	0	0
Other:	—	—	—	—	—	—	—	—	—	—

Counties Reporting Animal Rabies: Alexandria 1 fox, 6 raccoons; Arlington 1 raccoon; Augusta 1 cat; Fairfax 8 raccoons; Fauquier 1 skunk; Loudoun 1 skunk; Prince William 1 bat, 1 raccoon; Rockingham 1 skunk; Spotsylvania 1 skunk, 1 raccoon; Stafford 1 skunk, 1 raccoon

Occupational Illnesses: Occupational hearing loss 6; occupational pneumoconiosis 10; Asbestosis 3.

*3 year means

Published Monthly by the
VIRGINIA HEALTH DEPARTMENT
 Division of Epidemiology
 109 Governor Street
 Richmond, Virginia 23219

Bulk Rate
 U.S. POSTAGE
PAID
 Richmond, Va.
 Permit No. 729