




REPORTABLE DISEASE SURVEILLANCE IN VIRGINIA 2023



Office of Epidemiology
Virginia Department of Health

Introduction	3
Amebiasis	4
Anthrax.....	5
Arboviral Infection – Dengue.....	6
Arboviral Infection – Other	7
Arboviral Infection – West Nile Virus.....	8
Arboviral Infection – Zika Virus	9
Babesiosis.....	10
Botulism	11
Brucellosis.....	12
Campylobacteriosis.....	13
Chancroid.....	14
Chickenpox (Varicella)	15
<i>Chlamydia trachomatis</i> infection	16
Cholera	17
Cryptosporidiosis	18
Cyclosporiasis	19
Diphtheria.....	20
Ehrlichiosis/Anaplasmosis	21
Giardiasis	22
Gonorrhea	23
Granuloma inguinale.....	24
<i>Haemophilus influenzae</i> , invasive.....	25
Hantavirus Pulmonary Syndrome	26
Hemolytic Uremic Syndrome	27
Hepatitis A.....	28
Hepatitis B, acute.....	29
Hepatitis B, chronic	30
Hepatitis B, perinatal infection	31
Hepatitis C, acute.....	32
Hepatitis C, chronic.....	33
Hepatitis C, perinatal infection	34
Human immunodeficiency virus (HIV).....	35
Influenza A, novel virus	36
Influenza-associated deaths (less than 18 years of age).....	37
Legionellosis	38
Leprosy/Hansen’s Disease	39
Leptospirosis.....	40
Listeriosis	41
Lyme disease	42
Lymphogranuloma venereum	43
Malaria	44
Measles (Rubeola).....	45
Meningococcal disease (<i>Neisseria meningitidis</i>)	46
Mpox	47
Mumps	48
Paratyphoid infection	49
Pertussis	50
Plague.....	51

Poliovirus infection	52
Psittacosis	53
Q fever	54
Rabies, human	55
Rubella, including congenital rubella syndrome	56
Salmonellosis	57
Shiga toxin-producing <i>Escherichia coli</i> infection (STEC)	58
Shigellosis	59
Smallpox	60
Spotted Fever Rickettsiosis (including RMSF)	61
Streptococcal disease, Group A, invasive or toxic shock	62
<i>Streptococcus pneumoniae</i> , invasive (age less than 5 years)	63
Syphilis, early	64
Tetanus	65
Trichinosis (Trichinellosis)	66
Tuberculosis	67
Tularemia	68
Typhoid infection	69
Vaccinia adverse event following vaccination	70
Vancomycin-intermediate <i>Staphylococcus aureus</i> (VISA)	71
Vancomycin-resistant <i>Staphylococcus aureus</i> (VRSA)	72
Vibriosis, non-cholera	73
Viral hemorrhagic fever	74
Yellow fever	75
Yersiniosis	76

Introduction

The Virginia Department of Health's Office of Epidemiology is pleased to present its 2023 annual report of disease surveillance activities. The Office of Epidemiology, in partnership with health departments throughout Virginia, is responsible for the statewide surveillance of reportable diseases. Disease surveillance is the collection, analysis, interpretation, and dissemination of health-related data that is used to inform public health actions to reduce illness. This report summarizes surveillance data for diseases and conditions listed as reportable in the Virginia Regulations for Disease Reporting and Control and reported to the Centers for Disease Control and Prevention (CDC) each year.


Included in this report is a brief description of each disease, a map of disease incidence rate by locality or region, and a graph of reported cases each year for up to 10 years and the 5-year averages. Additional visualizations show disease data by age group, sex, and race for the year. For more information on disease reporting in Virginia, see the [Disease Reporting and Control Regulations](#) webpage.

Surveillance Data Notes


- The [Geography Locator Tool](#) can be used to identify Virginia's localities, health districts, and health planning regions.
- Conditions with an annual case count fewer than 20 are mapped at the regional level.
- If available data is insufficient to calculate a 5-year average, the 5-year average is not shown.
- Population data have been adjusted to represent the specific population under surveillance when data are restricted to certain age groups, including invasive *Streptococcus pneumoniae* infection (less than 5 years of age).
- Beginning in 2022, [Single Race Population Estimates from United States Census Bureau](#) using the prior year's population (e.g., 2023 surveillance data use 2022 Single Race population data) are used to calculate incidence rates.
- Patient Sex is reported as Female, Male, or Unknown as reported to VDH.
- Use of "Tiered Race," which merges race and ethnicity data, began in 2022. It is presented as Asian or Pacific Islander, Black, White, Hispanic, Native American, Two or More Races, Other, and Unknown.
- Report date reflects the CDC MMWR year (i.e., the date the case was reported to the health department). This may be different than the date of illness onset or diagnosis.
- Rates by locality are calculated based on residence of the patient, when known. When the address of the patient is neither reported by the healthcare provider nor ascertained by the health department, the location of the reporting source, such as the physician, hospital, or laboratory, is used.

Amebiasis

Amebiasis is an intestinal illness caused by the parasite *Entamoeba histolytica*. The condition is most common in tropical countries with poor sanitary conditions. In the United States, it is mainly identified in people who travel to or emigrate from these areas. Amebiasis can be spread through food or water that has been contaminated with feces from an infected person. Infected people are the only sources of the parasite. A person exposed to this parasite might experience mild or severe symptoms or no symptoms at all. Symptoms of amebiasis include diarrhea, stomach pain, and stomach cramping.




During 2023, 21 amebiasis cases were reported in Virginia. This represents a 22% decrease from the 27 cases reported in 2022. Sixty-two percent of cases reported their sex as male, and the statewide incidence rate was 0.2 per 100,000.




Anthrax

Anthrax is a serious infectious disease caused by the bacterium *Bacillus anthracis*. The bacteria occur naturally in soil and commonly affect domestic and wild animals around the world. People can get sick with anthrax if they come in contact with infected animals or contaminated animal products. People may be exposed to anthrax through inhalation, consuming contaminated food, or getting the bacteria on their skin. Depending on how a person was exposed, symptoms might include shortness of breath, sore throat, diarrhea, vomiting, headache, or skin sores. Anthrax is rare in the United States. However, sporadic outbreaks do occur in wild and domestic grazing animals, such as cattle or deer. To prevent anthrax exposures while visiting anthrax-endemic countries, travelers should avoid contact with animal carcasses and should not eat meat from animals butchered after having been found dead or ill.




No cases of anthrax were reported in Virginia in 2023.




Arboviral Infection – Dengue

Dengue is an arboviral illness that is transmitted by the bites of infected *Aedes aegypti* and *Aedes albopictus* mosquitoes. Cases are typically imported from tropical and subtropical countries, but transmission is increasingly common in certain parts of the United States, including Florida and Texas. There are four closely related virus types: Dengue-1, Dengue-2, Dengue-3, and Dengue-4. While most individuals will have an asymptomatic infection, approximately one in four will develop symptoms. Five percent of those infected experience severe dengue, a potentially fatal illness. Symptoms of dengue include fever, nausea/vomiting, rash, and aches and pains. Severe dengue may present with severe bleeding, plasma leakage, and organ involvement.




In 2023, Virginia reported 28 cases of dengue, all of which were imported. Cases were associated with travel to the Caribbean (8), South Asia (7), South America (8), and Central America (5). There has been an increase in cases when compared to the previous 5-year average of 9.8 cases. There were no reported fatalities or severe dengue illness in Virginia in 2023.




Arboviral Infection – Other

This category of arboviral infections is comprised of arboviral illnesses other than those caused by West Nile virus, Dengue, and Zika, and includes both endemic and non-endemic conditions. Arboviral infections are illnesses caused by arboviruses and are primarily transmitted to people by the bite of an infected mosquito, tick, or sandfly. Globally, there are over one hundred species of arboviruses that can cause disease in people. Due to ecologic, climatic, and cultural/economic factors, transmission of arboviral diseases have historically been limited in Virginia. Novel arboviral illnesses could emerge as ecological conditions become more favorable for non-native vectors and reservoirs.




In 2023, Virginia reported four cases of Chikungunya virus infection, all of which were acquired from international travel. Two cases of La Crosse virus infection were also reported; both cases were locally acquired and occurred among children.




Arboviral Infection – West Nile Virus

West Nile virus disease (WNV) is an arboviral illness that is transmitted by the bite of an infected mosquito. In Virginia, the primary vector is the mosquito *Culex pipiens*, also known as the “common house mosquito.” Transmission typically occurs during the summer and into the fall when the mosquito is most active. Most infected people will not have any symptoms, but one in five will become ill with symptoms that include fever, headache, body ache, nausea/vomiting, and rash. Approximately one out of every 150 infected people develop neuroinvasive infection with severe symptoms including high fever, disorientation, altered mental status, muscle weakness, and paralysis. Neuroinvasive infection may result in long term disability, and one in ten patients with neuroinvasive WNV will die from the illness.




Twelve cases of WNV were reported in Virginia in 2023; this was twice as many cases as reported in 2022. Nine cases were neuroinvasive, and three were non-neuroinvasive.




Arboviral Infection – Zika Virus

Zika virus disease (Zika) is a relatively new arbovirus of public health concern. Although it was discovered in 1948, transmission was extremely rare, and/or isolated geographically, until the epidemic of 2015-2016. Since 2017, local transmission of Zika has occurred in 89 countries globally. Zika is primarily transmitted by the bite of infected *Aedes* mosquitoes, but it can also spread from one person to another through sex or during pregnancy from mother to fetus. Many people infected with the virus will not have symptoms or will have only mild symptoms. The most common symptoms of Zika include fever, rash, conjunctivitis (red eyes), muscle and joint pains, and headache. Congenitally acquired infection is much more severe; symptoms in infants include central nervous system birth defects and smaller than expected head size (microcephaly).




Virginia did not report any cases of Zika virus in 2023. The last reported case in a Virginia resident occurred in 2021, which was acquired during travel to Mexico.




Babesiosis

Babesiosis is a disease caused by the protozoa, *Babesia*. There are two species that cause illness in humans in the U.S., including *Babesia microti* and *Babesia duncani*, with the former causing more than ninety-nine percent of human cases. Babesiosis is most often acquired from the bite of infected blacklegged ticks, but blood transfusion and transplacental/perinatal transmission are also possible. Symptoms can begin up to two months after a tick bite and range from flu-like symptoms (fever, chills, headache, myalgia) to life-threatening conditions, such as disseminated intravascular coagulation, acute respiratory distress, and myocardial infarction. Illness usually occurs from late spring to early fall. The elderly and immunocompromised are most susceptible.




Eleven cases of babesiosis were reported in Virginia residents in 2023. All cases were acquired by tickborne transmission. Seven of these cases were acquired in Virginia. The Eastern Shore had three locally acquired cases.




Botulism

Botulism is a rare but serious illness caused by a toxin produced by the spore-forming bacterium, *Clostridium botulinum*. The bacterium that makes the toxin is found in the soil, and sometimes in water. The disease is classified as foodborne when resulting from ingestion of food contaminated with the toxin, or non-foodborne (infant or wound botulism) when resulting from ingestion of food contaminated with spores that form toxin in the intestine, or contamination of an open wound with soil containing spores. Symptoms of botulism include double vision, blurred vision, drooping eyelids, slurred speech, difficulty swallowing, and muscle weakness. Infants with botulism appear very tired, feed poorly, are constipated, and have a weak cry and poor muscle tone.




Five cases of botulism were reported in Virginia in 2023. All cases reported their sex as female, and all were younger than one year of age and were classified as non-foodborne infant botulism. The statewide incidence rate was 0.1 per 100,000 population.



Brucellosis

Brucellosis is a disease caused by the bacterium, *Brucella*. The bacterium is most commonly transmitted through eating or drinking unpasteurized milk or milk products from infected animals. It can also be transmitted by contamination of skin wounds with infected animal tissue or body fluids and by inhalation of the organism. Symptoms include intermittent or irregular fever, headache, chills, sweating, and muscle pain. Some signs and symptoms may persist for longer periods of time, while others may never resolve. The best way to prevent brucellosis is by not consuming unpasteurized dairy products and by using barrier precautions, such as wearing rubber gloves, when handling animal tissue. Laboratory workers should take proper infection control precautions when handling samples that might contain *Brucella*.




No cases of brucellosis were reported in 2023. One case of brucellosis was reported in 2022. Prior to 2022, the last time Virginia reported a case of brucellosis was in 2020 when two cases were reported. Virginia typically reports fewer than five cases of brucellosis each year.


Age Group, 2023								Sex, 2023			Tiered Race, 2023							
											00000000							
0	0	0	0	0	0	0	0				Asian or Pacific Islander	Black	White	Hispanic	Native American	Other	Unknown	Two or More Races
<1	1-9	10-19	20-29	30-39	40-49	50-59	60+											

Campylobacteriosis

Campylobacteriosis is an infection caused by the bacteria *Campylobacter*, which affects the intestinal tract and causes diarrhea. The bacteria are commonly found in the gut of animals and birds, which carry the bacteria without becoming ill. It can be transmitted to people by drinking unpasteurized (raw) milk or dairy products or contaminated water, or eating undercooked meats, especially chicken. Symptoms of campylobacteriosis can include mild to severe diarrhea, fever, and stomach cramping. Campylobacteriosis is one of the most common causes of diarrheal illness in the United States. Transmission of this condition can be reduced through proper hand washing, consuming pasteurized dairy products, and ensuring poultry is cooked to appropriate temperatures.




In 2023, 1,749 campylobacteriosis cases were reported in Virginia, with a statewide incidence rate of 20.2 per 100,000 population. Adults 60 years of age and older represented 31% of all cases.



Chancroid

Chancroid is caused by the bacterium *Haemophilus ducreyi*. It is transmitted through sexual skin-to-skin contact with open sores. Symptoms usually occur five to ten days after exposure. Chancroid is characterized by multiple lesions or sores on the genitals which become purulent and eventually rupture. Additional symptoms include inguinal adenopathy (swollen lymph nodes), painful urination, vaginal discharge, rectal bleeding, pain with bowel movements, and dyspareunia. Chancroid became a nationally notifiable condition in 1944, and the case definition has not changed since 1996. Infections of Chancroid might still occur in certain African regions and the Caribbean, but are rare in other parts of the world. Since 2012, there have only been 70 reported cases of chancroid in the US.




In 2023, there were no reported cases of chancroid in Virginia.


Age Group, 2023									Sex, 2023				Tiered Race, 2023							
0	0	0	0	0	0	0	0	0					0	0	0	0	0	0	0	0
<1	1-9	10-19	20-29	30-39	40-49	50-59	60+						Asian or Pacific Islander	Black	White	Hispanic	Native American	Other	Unknown	Two or More Races
									Case Count	%	Annual Rate									
									Female	0	0.0									
									Male	0	0.0									
									Unknown	0	0.0									

Chickenpox (Varicella)

Chickenpox is a highly contagious disease caused by the varicella-zoster virus (VZV). The virus can be spread in the air when a person coughs or sneezes, or by direct contact with either the chickenpox or shingles rash before a scab forms. Initial symptoms include sudden onset of fever, headache, and fatigue. An itchy, blister-like rash, usually starting on the face, chest, or back, follows 1-2 days later. In the United States, most cases occur in young, school-aged children. The risk of chickenpox is low in persons who have received two doses of vaccine.




Virginia reported 285 cases of chickenpox in 2023, which is nearly double the number of cases reported in 2022. This increase was primarily due to outbreaks in K-12 schools, universities, daycares, and correctional facilities.




Chlamydia trachomatis infection

Chlamydia is a common sexually transmitted disease caused by *Chlamydia trachomatis*. It is the most frequently reported bacterial infection in the United States, particularly among women and young people ages 15-24. Anyone can get chlamydia through vaginal, oral, or anal sex with someone who has the infection. It may also be transmitted to infants from the genital tract of an infected mother during delivery. Asymptomatic infection is common, so many people are unaware of their infection. Symptoms may include vaginal, penile, or rectal discharge and/or burning during urination. Chlamydial infection in women can result in serious sequelae, including pelvic inflammatory disease (PID), ectopic pregnancy, and infertility. Transmission of this disease can be mitigated by regularly screening and prompt treatment for sexually active individuals.




In 2023, 41,206 chlamydia cases were reported in Virginia. The rate of reported chlamydia in 2023 (474.8 per 100,000) was similar to the rate in 2022 (471.1 per 100,000). From 2022 to 2023, the rate of reported chlamydia increased among men (2%) and remained the same among women, yet cases in women accounted for 64% of all reported chlamydia in 2023. The rate of chlamydia was highest among persons aged 20 to 29 years (1,971.5 per 100,000) and 55% of cases were reported among this age group.



Cholera

Cholera is an acute diarrheal illness caused by certain strains of the bacterium *Vibrio cholerae*. People with cholera shed the bacteria in their feces (stool). It is transmitted when feces from a person with cholera gets into the water supply and people drink the contaminated water. Food can also become contaminated if it is washed with unclean water or fertilized with sewage or contaminated soil. Shellfish harvested from water that has been contaminated can cause people to get sick if the shellfish are eaten raw or undercooked. Symptoms can range from mild to severe and commonly include profuse watery diarrhea, vomiting, and weakness. Cholera is not a threat in the United States and other countries with advanced water and sanitation systems.




No cases of cholera were reported in Virginia in 2023. The last reported case of cholera occurred in 2022 and was associated with international travel.


Age Group, 2023									Sex, 2023				Tiered Race, 2023							
0	0	0	0	0	0	0	0	0	0 0 0 0 0 0 0 0											
<1	1-9	10-19	20-29	30-39	40-49	50-59	60+													
									Case Count % Annual Rate											
									Female 0 0.0				Asian or Pacific Islander							
									Male 0 0.0				Black							
									Unknown 0 0.0				White							
													Hispanic							
													Native American							
													Other							
													Unknown							
													Two or More Races							

Cryptosporidiosis

Cryptosporidiosis is a diarrheal disease caused by the parasite *Cryptosporidium*. The parasite is found in the intestines of infected humans and animals and is shed in feces. People become infected by eating food or drinking water contaminated with the parasite, or by swimming in contaminated lakes, streams, rivers, or swimming pools. The parasite is protected by an outer shell and is very resistant to chlorine disinfection. Cryptosporidiosis causes watery diarrhea and abdominal cramping, and sometimes vomiting and low-grade fever. Transmission of cryptosporidiosis can be reduced through proper hand washing, not using public swimming facilities while ill with diarrhea, and avoiding drinking water directly from streams, lakes, springs, or any unknown water source.




During 2023, 283 cases of cryptosporidiosis were reported in Virginia with a statewide incidence rate of 3.3 cases per 100,000 population. Fifty-four percent of cases reported their sex as female.




Cyclosporiasis

Cyclosporiasis is an infection of the intestine caused by the parasite *Cyclospora*. Cyclosporiasis is transmitted through feces and people become infected by eating food or drinking water contaminated with the parasite. The infection can occur in many countries, including the United States, but it is more common in tropical areas. Outbreaks of cyclosporiasis in the United States have been linked to imported fresh produce, such as berries, lettuce, and fresh herbs. Most cases and outbreaks are reported during the summer months, although infections can occur throughout the year. Cyclosporiasis can cause watery diarrhea, stomach cramps, loss of appetite, and weight loss, and some people who are infected do not have any symptoms.




In 2023, 183 cases of cyclosporiasis were reported in Virginia with a statewide incidence rate of 2.1 per 100,000 population. This increase was attributed to a cluster of domestically acquired cases identified during the summer months of 2023, which included 159 cases. Unfortunately, a definitive food vehicle was not identified for this cluster. Fifty-four percent of cases reported their sex as female.




Diphtheria

Diphtheria is a serious disease caused by a toxin produced by the bacterium *Corynebacterium diphtheriae*. Diphtheria can infect the nose, throat, or skin, and can be deadly. Signs and symptoms include weakness, sore throat, mild fever, swollen glands in the neck, and open sores or ulcers. The overall case-fatality rate for diphtheria is 5-10%, with higher death rates (up to 20%) among persons younger than 5 and older than 40 years of age. In the U.S., cases gradually declined after the vaccine became routinely used in the 1940s, and cases remain rare today.




Ehrlichiosis/Anaplasmosis

Ehrlichiosis and anaplasmosis are bacterial diseases transmitted by the bite of an infected tick. The four subtypes which cause illness are *Ehrlichia chaffeensis*, *Ehrlichia ewingii*, *Anaplasma phagocytophilum* and Undetermined human ehrlichiosis/anaplasmosis. Both *E. chaffeensis* and *E. ewingii* are transmitted by the lone star tick, while *A. phagocytophilum* is transmitted by the blacklegged tick. Symptoms include fever, headache, body aches, anemia, low white blood cell count, low platelet count, and elevated liver enzymes. Severe cases can present with organ failure, altered mental status, and encephalopathy. In general, ehrlichiosis cases are more severe than anaplasmosis cases. The best way to prevent this infection is to limit tick exposure.




There were 98 cases of *E. chaffeensis* infection, 31 cases of *A. phagocytophilum* infection, and three cases of ehrlichiosis/anaplasmosis undetermined infection in Virginia in 2023. There was one reported fatality caused by *E. chaffeensis* infection. The combined total (132) is higher than the previous 5-year average of 121 cases, but within the typical number of cases reported per year for the last ten years.




Giardiasis

Giardiasis is a diarrheal illness caused by the microscopic parasite *Giardia*. The parasite can be found in the feces (stool) of infected people and animals, which can further contaminate soil, food, water, or any surface which comes in contact with the feces. People get sick by swallowing contaminated food or water or by putting their hands in their mouths after touching contaminated surfaces. Anyone can get giardiasis, but it tends to occur more often in international travelers, and persons who drink improperly treated surface water (such as hikers drinking from a stream or people swallowing water while swimming in a river or lake). Transmission of giardiasis can be reduced through proper hand washing, and avoiding drinking water directly from streams, lakes, springs, or any unknown water source.




In 2023, 307 cases of giardiasis were reported in Virginia with a statewide incidence rate of 3.5 per 100,000 population. Sixty-seven percent of giardiasis cases reported their sex as male.




Gonorrhea

Gonorrhea, caused by the bacterium *Neisseria gonorrhoeae*, is the second most commonly reported bacterial disease. It is primarily transmitted through sexual contact but can also be passed perinatally from mother to baby during childbirth. Many infections are asymptomatic; however, symptoms may occur within 7 to 30 days of exposure. Symptoms include painful urination, penile/vaginal discharge, intermenstrual bleeding, painful bowel movements, and dyspareunia. Among women, untreated gonococcal infections can result in pelvic inflammatory disease (PID), tubal scarring, and infertility or ectopic pregnancy. Gonorrhea is curable, but proper antibiotic treatment has become more difficult over time due to increases in drug-resistant strains. Populations at a higher risk for gonorrhea include males, non-Hispanic black persons, and those 20-39 years of age.




In 2023, 13,795 gonorrhea cases were reported in Virginia, representing a 2% increase in incidence rate from 2022 (155.2 to 158.9 per 100,000). Rates continue to be highest among males (194.0 per 100,000) and Black persons (435.5 per 100,000); rates increased in both subgroups from 2022 (7% and 11% increase, respectively). The rate of reported gonorrhea was highest among persons aged 20 to 29 years in 2023 (564.7 per 100,000).




Granuloma inguinale

Granuloma inguinale (donovanosis) is a chronic, relapsing, granulomatous anogenital infection caused by the bacterium *Klebsiella granulomatis* (formerly known as *Calymmatobacterium granulomatis*). Granuloma inguinale occurs sporadically in some tropical and developing countries but is rare in other parts of the world. It can be transmitted through sexual contact with open sores. Usually, symptoms occur one to twelve weeks after exposure. Clinically, the disease is characterized by painless, slowly progressive nodules that break down into shallow, sharply demarcated ulcers. Lesions may occur on the skin, genitalia, or perineal areas and slowly spread to the lower abdomen and thighs. Treatment has been reported to halt the progression of lesions. Granuloma inguinale became nationally notifiable in 1942, and the case definition has remained the same since 1990.




In 2023, there were no reported cases of granuloma inguinale in Virginia.




Haemophilus influenzae, invasive

Haemophilus influenzae is a type of bacteria commonly found in the nose and throat. The bacteria are transmitted through respiratory droplets and can invade the body to cause serious infections, including pneumonia and meningitis. Symptoms of meningitis may include fever, headache, neck stiffness, vomiting, and nausea. Pneumonia symptoms may include shortness of breath, fever, chest pain, and cough. *H. influenzae* type B (Hib) was once one of the leading causes of bacterial meningitis among young children. Since 1991, cases in children under 5 years of age have decreased 99% due to widespread use of the Hib vaccine.




Virginia reported 198 cases of invasive *H. influenzae* in 2023, a 24% increase from the year prior, with the greatest burden in the Eastern and Southwest regions.



Hantavirus Pulmonary Syndrome

Hantaviruses are a family of viruses that are mainly spread by rodents and can cause varied disease syndromes in people worldwide. Infection with any hantavirus can produce hantavirus disease in people. Hantaviruses in the Americas are known as “New World” hantaviruses and may cause hantavirus pulmonary syndrome (HPS). Early symptoms include fever and muscle pain, gastrointestinal complaints, headaches, and dizziness. These symptoms may be followed by an abrupt onset of respiratory distress and decreased blood pressure, which can be fatal. Hantavirus infections can be prevented by excluding rodents from houses and other buildings. Protective measures include disinfecting rodent-contaminated areas with a spray disinfectant solution prior to cleaning. Contaminated areas should be cleaned with a wet mop and not be vacuumed or swept to decrease the likelihood of creating an aerosol containing the virus.




No cases of hantavirus pulmonary syndrome (HPS) were reported in Virginia in 2023. The last case of HPS reported in Virginia was in 2021.


Age Group, 2023								Sex, 2023			Tiered Race, 2023							
0	0	0	0	0	0	0	0	0			0	0	0	0	0	0	0	0
<1	1-9	10-19	20-29	30-39	40-49	50-59	60+				Asian or Pacific Islander	Black	White	Hispanic	Native American	Other	Unknown	Two or More Races
								Case Count	%	Annual Rate								
								Female	0	0.0								
								Male	0	0.0								
								Unknown	0	0.0								

Hemolytic Uremic Syndrome

Hemolytic Uremic Syndrome (HUS) is a serious disease that affects the kidneys and blood clotting system, which can cause the sudden development of kidney failure and damage to other organs. Most cases of HUS occur as a rare complication of infection with a diarrheal illness caused by a toxin-producing bacterium such as *E. coli*. People become infected with *E. coli* by touching contaminated surfaces and then putting their hands in their mouths, being in contact with cattle and other farm animals, or by eating or drinking contaminated food or water. Early symptoms of HUS include decreased urine output, diarrhea, and feeling slow and tired (lethargy). Anyone can get HUS, but it is more common in children than adults, especially children less than five years of age.




During 2023, 7 cases of HUS were reported in Virginia with a statewide incidence rate of 0.1 per 100,000 population. Two HUS cases were associated with Shiga toxin-producing *E. Coli* (STEC) outbreaks. Children ages one to nine represented 71% of all cases reported.




Hepatitis A

Hepatitis A is a contagious liver disease caused by infection with the hepatitis A virus (HAV). Hepatitis A ranges in severity from a mild disease lasting several weeks to a severe disease lasting several months. HAV is found in the feces of infected people, whose feces can contaminate surfaces, objects, food, or water. HAV is spread by consuming contaminated food or water, or through person to person contact through sexual activities and sharing contaminated needles. An increase in hepatitis A in many states across the country was identified in 2016 among adults of high-risk groups. Virginia was considered an "outbreak state" in 2019 and has been experiencing an increased number of cases statewide since. The best way to prevent hepatitis A is to get vaccinated, and wash hands with soap and water after using the bathroom, changing diapers, and before preparing and eating food.




In 2023, 34 hepatitis A virus cases were reported in Virginia with a statewide incidence rate of 0.4 per 100,000 population. Young adults ages 20 to 29 represented 35% of cases and 55% of cases reported their sex as female.




Hepatitis B, acute

Hepatitis B virus (HBV) is a hepadnavirus that can cause acute (short-term) or chronic (long-term) infection. The acute hepatitis B case definition was revised in 2012. This report represents newly identified probable and confirmed acute hepatitis B cases in Virginia.




In 2023, 31 newly identified acute hepatitis B cases were reported; a 28% decrease compared to 2022. In 2022, 43 acute hepatitis B cases were reported; a 9% increase compared to 2021. No outbreaks associated with HBV were reported in 2023.




Hepatitis B, chronic

Hepatitis B virus (HBV) is a hepadnavirus that can cause acute (short-term) or chronic (long-term) infection. The chronic hepatitis B case definition was revised in 2012. This report represents newly identified probable and confirmed chronic hepatitis B cases in Virginia.




In 2023, 1,803 newly identified chronic hepatitis B cases were reported, representing a 7% increase compared to 2022, signaling minimal change in disease burden compared to the prior year. In 2022, 1,684 cases were reported which represented a 2% decrease compared to 2021. No outbreaks with HBV were reported in 2023.




Hepatitis B, perinatal infection

Perinatal hepatitis B (HBV) transmission can occur at birth when a pregnant HBV-infected woman delivers her infant. Transmission is preventable by identifying HBV-infected pregnant women and providing post exposure prophylaxis (PEP), consisting of hepatitis B immune globulin (HBIG) and hepatitis B (HepB) vaccine to their infants within 12 hours of birth. Without administration of PEP, at birth, approximately 40% of infants born to HBV-infected mothers in the United States will develop chronic HBV infection. 0.7%-1.1% of infants will develop infection when PEP is given within 12 hours of birth and the HepB vaccine series is completed. Infants at greatest risk to infection, even when PEP is given, are born to mothers with high viral loads. In Virginia, an average of 217 infants are identified annually as being born to HBV-infected women.




In 2023, no cases of perinatal hepatitis B were identified. The Virginia Perinatal Hepatitis B Prevention Program conducted follow up on 475 infants born between 2019 and 2023. Of those, 204 infants were determined to not be infected with hepatitis B, while 271 infants continued to be monitored into 2024.




Hepatitis C, acute

Hepatitis C virus (HCV) is the most common blood-borne infection in the United States and can cause acute (short-term) or chronic (long-term) infection. The acute hepatitis C case definition was last updated in 2020. The statistics in this report represent newly identified probable and confirmed acute hepatitis C cases. The number of newly identified cases increased substantially from 41 in 2013 to 98 in 2019.




In 2023, 36 newly identified acute hepatitis C cases were reported, a 13% increase compared to 2022. In 2022, 32 acute hepatitis C cases were reported, an 18% decrease compared to 2021. No outbreaks associated with HCV were reported in 2023.




Hepatitis C, chronic

Hepatitis C virus (HCV) is the most common blood-borne infection in the United States and can cause acute (short-term) or chronic (long-term) infection. The chronic hepatitis C case definition was last updated in 2020. The statistics included in this report represent newly identified probable and confirmed chronic hepatitis C cases. The number of newly identified cases increased substantially from 6,012 in 2013 to 11,274 in 2019.




In 2023, 4,439 newly identified chronic hepatitis C cases were reported, a 23% decrease compared to 2022. In 2022, 5,772 chronic hepatitis C cases were reported, a 11% decrease compared to 2021. No outbreaks associated with HCV were reported in 2023.




Hepatitis C, perinatal infection

Hepatitis C virus (HCV) is the most common blood-borne infection in the United States and can cause acute (short-term) or chronic (long-term) infection. Perinatal hepatitis C became reportable in Virginia in 2018. The statistics included in this report represent newly identified confirmed perinatal hepatitis C cases. Infants with a positive HCV RNA nucleic acid test or with a detectable HCV genotype at ≥ 2 months and ≤ 36 months of age, and not known to have been exposed to HCV via a mechanism other than perinatal exposure, are classified as perinatal hepatitis C cases.




In 2023, 11 perinatal hepatitis C cases were reported, a 38% increase compared to 2022. Eight perinatal hepatitis C cases were reported each year in 2022 and 2021. No outbreaks associated with HCV were reported in 2023.




Human immunodeficiency virus (HIV)

Human immunodeficiency virus (HIV) is a preventable virus that attacks the body's immune system and can lead to acquired immunodeficiency syndrome (AIDS) if left untreated. There is no cure for HIV, but in people who take antiretroviral treatment (ART) as prescribed by their physician, the virus can become undetectable, making them unable to pass the virus to others.




From 2021 to 2023, new HIV diagnoses in Virginia increased. In 2023, there were 865 new HIV diagnoses in Virginia. Among new diagnoses in 2023, people aged 30-34 years had the highest number of new cases, followed closely by people aged 20-24 and 25-29. More men (n=721, 83%) than women (n=144, 16%) were newly diagnosed (determined by sex at birth); and people who reported Black or African American race were more likely to be newly diagnosed with HIV than other races. Since 2020, the number of people who reported Hispanic or Latino ethnicity increased.



Influenza A, novel virus

A novel influenza virus is an influenza A virus with a subtype that is different from the flu viruses that usually spread in people (H3N2 and H1N1). Some examples include H7N9 and H5N1. Occasionally, strains of influenza that normally affect birds, pigs, and other animals can infect humans. When flu viruses that normally affect pigs (swine flu viruses) cause infections in humans, these viruses are called variant influenza viruses. Sometimes, human infections with novel or variant flu viruses occur because of the close contact between humans and animals. Other times, the infections occur because of changes in the influenza virus. Human infection with novel influenza A viruses became nationally notifiable in the U.S. in 2007, and the case definition was last updated in 2014. Novel influenza A viruses are of concern because of the potential impact they could have on public health if they gain the ability to spread easily from person to person. The best way to prevent infection is to avoid sources of exposure whenever possible.




For 2023, there were zero cases of novel influenza A virus reported. The Virginia prior 5-year average has remained at zero cases.


Age Group, 2023									Sex, 2023				Tiered Race, 2023							
0	0	0	0	0	0	0	0	0	0 0 0 0 0 0 0 0											
<1	1-9	10-19	20-29	30-39	40-49	50-59	60+													
									Case Count % Annual Rate											
									Female 0 0.0				Asian or Pacific Islander							
									Male 0 0.0				Black							
									Unknown 0 0.0				White							
													Hispanic							
													Native American							
													Other							
													Unknown							
													Two or More Races							

Influenza-associated deaths (less than 18 years of age)

Influenza (“flu”) is a contagious respiratory illness caused by influenza viruses that infect the nose, throat, and sometimes the lungs. Flu can cause mild to severe illness and at times can lead to death. For surveillance purposes, an influenza-associated pediatric death is defined as a death in a person <18 years of age, resulting from a clinically compatible illness that was confirmed to be flu, without a period of complete recovery between the illness and death. Virginia has been monitoring flu-associated pediatric deaths since 2003, with reporting required by Virginia Administrative Code in May 2007. Surveillance of flu deaths is one component used to measure the severity and impact of flu illness. The best way to reduce risk from seasonal flu and its potentially serious complications is to receive an annual flu vaccine.




Four flu-associated pediatric deaths were reported in Virginia in 2023. Of these, three occurred among the 5-12 years age group, and one occurred among the 13-17 years age group. On average, there are three flu-associated pediatric deaths reported each year in VA.




Legionellosis

Legionellosis is the collective name for respiratory infections caused by *Legionella* bacteria. *Legionella* bacteria commonly cause one of two lung diseases: a serious type of pneumonia (lung infection) called Legionnaires' disease, or Pontiac fever, which is a less severe respiratory disease. Rarely, *Legionella* can cause infections outside of the lungs, such as heart or wound infections. Antibiotics are used to treat *Legionella* infections. Most healthy people exposed to *Legionella* don't get sick. However, certain factors such as smoking status, older age, and health issues or conditions put some people at increased risk for getting Legionnaires' disease. The key to preventing Legionnaires' disease is to reduce the risk of *Legionella* growth and spread.




In Virginia, the 5-year average for legionellosis has been increasing steadily over the last 10 years. An increase in cases is also being reported nationwide. Of the 171 legionellosis cases reported in 2023, 109 (64%) were among people 60 years of age or older, with a rate of 5.4 cases per 100,000 population.




Leprosy/Hansen's Disease

Hansen's disease (also known as leprosy) is an infection caused by slow-growing bacteria called *Mycobacterium leprae*. It can affect the nerves, skin, eyes, and lining of the nose (nasal mucosa). It is not known exactly how Hansen's disease spreads between people. Scientists currently think it may happen when a person with Hansen's disease coughs or sneezes, and a healthy person breathes in the droplets containing the bacteria. However, prolonged, close contact with someone with untreated leprosy over many months is needed to catch the disease. In the southern United States, some armadillos are naturally infected with this bacterium. For general health reasons, avoid contact with armadillos whenever possible.




One case of Hansen's disease was reported in 2023 in Virginia, which was thought to be related to exposure in another country. Prior to 2023, the last case of Hansen's disease in Virginia was reported in 2019.




Leptospirosis

Leptospirosis is a disease that affects humans and animals and is caused by bacteria *Leptospira*. In humans, it can cause a wide range of symptoms, such as fever, muscle aches, headache, and vomiting, which may be mistaken for other diseases. Some infected persons, however, may have no symptoms at all. People may become infected through contact with urine of infected animals, or water, soil, or food contaminated with the urine of infected animals. The risk of acquiring leptospirosis can be greatly reduced by not swimming or wading in water that might be contaminated with animal urine or eliminating contact with potentially infected animals. Protective clothing or footwear should be worn by those exposed to contaminated water or soil because of their job or recreational activities.




Virginia reported two cases of leptospirosis in 2023 among patients whose primary risk was contact with a contaminated body of water. Virginia typically reports no more than three leptospirosis cases per year.




Listeriosis

Listeriosis is a serious disease usually caused by eating food contaminated with the bacterium *Listeria monocytogenes*. Illness develops when the bacteria get into sterile body sites. Listeriosis mainly affects pregnant women, newborns, adults aged 65 years or older, and people with weakened immune systems. Serious illness in people without these risk factors is rare. *Listeria* bacteria are commonly found in the environment, including in soil and water. Consuming contaminated foods (unwashed raw fruits and vegetables, raw milk, soft cheeses, deli meats) can cause infection. Pregnant women with listeriosis typically experience fever and mild flu-like symptoms such as fatigue and muscle aches. Older adults and people with weakened immune systems might experience fever, muscle aches, headache, stiff neck, confusion, loss of balance, and convulsions.




During 2023, 31 listeriosis cases were reported in Virginia with a statewide incidence rate of 0.4 per 100,000 population. Adults over 60 years of age represented 65% of listeriosis cases reported and 61% of cases reported their sex as male.




Lyme disease

Lyme disease is an illness caused by the bacterium *Borrelia burgdorferi*, and less commonly, *Borrelia mayonii*. It is transmitted by the bite of an infected blacklegged tick. Symptoms usually appear within 30 days of the tick bite, often beginning with an erythema migrans rash. If untreated, illness can present as joint pain, headache, fatigue, and swollen lymph nodes. Advanced, untreated illness can cause facial palsy, severe arthritis, irregular heartbeat, meningitis, and nerve pain. Some patients may develop the condition Post-Treatment Lyme Disease Syndrome and continue to have symptoms even after completing a prescribed antibiotic regimen. Children aged 1-9 years and those aged 60 years and older have the highest disease burden among age groups. In 2022, there was a change in the case definition which has caused a continued rise in Lyme disease cases from previous years.




In 2023, Virginia reported 1,747 cases of Lyme disease. The previous 5-year average was 1,000 cases. There was a continued rise in Lyme disease cases from previous years due to the new case definition.




Lymphogranuloma venereum

Lymphogranuloma venereum (LGV) is a sexually transmitted infection caused by the bacterium *Chlamydia trachomatis*, specifically serovars L1, L2, or L3. It can cause severe inflammation and invasive infection. Common presentations of LGV infection include proctocolitis (after rectal exposure) and inflammation of the lymph nodes in the genital area. Sometimes a self-limited genital ulcer or papule can occur at the site of inoculation. Some infections may be asymptomatic. LGV became nationally notifiable in 1941 but was removed as a separate notifiable condition in 1995 when chlamydia was added. However, in 2022 the case definition for chlamydia was updated to distinguish LGV from other infections due to chlamydia. LGV has historically been rare in developed counties, with only occasional outbreaks primarily occurring among men who have sex with men.




Malaria

Malaria is a mosquito-transmitted illness caused by several species of parasitic *Plasmodium*. The clinical presentation can range from flu-like to severe illness, and symptoms include fever, muscle pain, chills, sweats, headache, nausea, vomiting, anemia, and jaundice. The vast majority of cases reported in the United States are travel-related; endemic regions with ongoing transmission include Africa, Central/South America, and Southern Asia. However, locally acquired cases have occurred in the Commonwealth in both Westmoreland County in 1998 and Loudoun County in 2002. It is recommended that travelers to malaria-endemic regions take anti-malaria drugs; the specific drug regimen is determined by the area of travel.




There were 127 cases of malaria reported in Virginia in 2023. All cases were acquired outside of the United States and the majority of cases (55%) were acquired via travel to Sierra Leone (43), Ghana (19), and Cameroon (8).




Measles (Rubeola)

Measles (Rubeola) is a highly contagious respiratory illness spread by direct contact with infectious droplets or by airborne spread when an infected person breathes, coughs, or sneezes. Symptoms include fever and a rash that starts on the face and spreads to the neck, trunk, arms, and legs. A cough, runny nose, or red, itchy eyes might also be present. Measles cases declined significantly in the United States after the live measles vaccine was licensed in 1963, and measles was declared eliminated from the U.S. in 2000. Today, most measles cases are reported in the U.S. as a result of importations by unvaccinated people who were infected while in other countries.




One measles case associated with international travel was reported in Virginia in 2023.




Meningococcal disease (*Neisseria meningitidis*)

Meningococcal disease is a serious illness caused by the bacterium *Neisseria meningitidis*, which is spread from person to person through respiratory and throat secretions. Some people may develop a serious form of illness, such as meningitis (inflammation of the lining of the brain and spinal cord) or a bloodstream infection. Signs and symptoms include fever, headache, stiff neck, nausea, vomiting, photophobia, and altered mental status. It is most common in infants, children, adolescents, and young adults. Ensuring timely receipt of meningococcal vaccination is the best way to protect against meningococcal disease.




Virginia reported 26 cases of meningococcal disease in 2023, an increase of over 60% from 2022. The majority of cases (81%) were linked to a statewide outbreak of *Neisseria meningitidis* serogroup Y, first identified in Virginia's Eastern Region in June 2022. Groups disproportionately affected by the outbreak included Black or African American adults, people 30-60 years of age, and people with HIV.




Mpox

Mpox is a disease caused by infection with the monkeypox virus. It is spread through close contact with infected people or animals. People with mpox often present with a rash, which begins as painful or itchy blister-like lesions that scab over as they heal. Other symptoms include fever, chills, swollen lymph nodes, fatigue, aches, and respiratory symptoms such as a sore throat or cough. Illness severity can range from mild to severe. A vaccine is available and is recommended for persons at risk. Mpox was reportable in Virginia from 2003-2016, during which no cases were reported. Mpox again became reportable in Virginia in 2022 in response to a worldwide outbreak of mpox.




In 2023, 12 cases of mpox were reported in Virginia. All of the cases were reported in the Northern region. The first mpox case in Virginia was reported in May 2022, with cases peaking in August 2022.




Mumps

Mumps is a contagious disease caused by a paramyxovirus. It can be transmitted person to person through direct contact with saliva or respiratory droplets of a person infected with mumps. Signs and symptoms include fever, headache, muscle aches, fatigue, and swelling of salivary glands under the ears on one or both sides (parotitis). Significant outbreaks have happened in settings where people have intense or frequent close contact (e.g., college campuses).




Virginia reported 14 mumps cases in 2023, a slight decrease from the year prior.




Paratyphoid infection

Paratyphoid infection is a disease caused by the bacterium *Salmonella Paratyphi*. Anyone can get paratyphoid infection, but it occurs more often in people who travel to developing countries in Asia, Africa, and South and Central America, where the disease is common. It can be transmitted by eating or drinking food or water that has been contaminated with feces or urine of people with the disease or by direct contact with a person who has the disease. Paratyphoid infection can cause high fever, headaches, weakness, loss of appetite, and diarrhea, or constipation. When traveling, particularly to areas where paratyphoid infection is common, it is important to practice safe food and water habits. These include eating food that has been fully cooked, drinking water (and ice) from a safe source, avoiding raw or undercooked food, and avoiding tap or well water (or ice made with tap or well water).




In 2023, 9 paratyphoid infection cases were reported in Virginia, with a statewide incidence rate of 0.1 per 100,000 population. Sixty-seven percent of cases reported their sex as female.




Pertussis

Pertussis (whooping cough) is a very contagious respiratory disease caused by the bacteria *Bordetella pertussis*. These bacteria spread through the air in droplets produced when someone sneezes or coughs. The illness begins with cold-like symptoms, including runny nose, mild fever, and cough. The cough lasts 1-2 weeks, gradually worsens, and may include coughing fits, leading to a high-pitched "whoop," vomiting, or struggling to breathe. It most commonly occurs in very young children who have not been vaccinated.




Virginia reported 120 cases of pertussis in 2023, an increase of 64% from the previous year. Reasons for this change include an increase in the use of PCR testing for pertussis diagnosis and a return to more typical pertussis trends following the COVID-19 pandemic.




Plague

Plague is caused by the bacteria *Yersinia pestis*. It can be transmitted to humans through the bite of infected fleas or through handling the tissue or body fluids of a plague-infected animal. If the disease attacks the lungs, it can be spread from person-to-person by respiratory droplets released when coughing. Clinical symptoms in people include fever, chills, nausea, headache, and body aches. Other clinical presentations include swollen lymph nodes (“buboes”), bloodstream infections, and pneumonia. In areas where plague occurs, travelers should avoid contact with rodents and fleas and avoid handling stray animals. Presently, human plague infections occur rarely in rural areas in the western United States. Most human cases since the 1990s have occurred in Africa. Modern antibiotics are effective in treating plague.



Poliovirus infection

Polio, or poliomyelitis, is a disabling and life-threatening disease caused by the poliovirus. Poliovirus is very contagious and spreads through contact with the feces of an infected person or contact with droplets from a sneeze or cough of an infected person (less common). Most people who are infected with poliovirus will not have any visible symptoms; however, a small portion of people will develop serious symptoms that affect the brain and spinal cord, including meningitis or paralysis. Following the introduction of inactivated polio vaccine (IPV) in 1955 and oral polio vaccine (OPV) in 1961, the reported incidence of poliomyelitis in the United States declined dramatically, with under 100 cases in 1965 and to under 10 cases in 1973.




The last reported cases of poliomyelitis in Virginia occurred in 1978.

Age Group, 2023								Sex, 2023			Tiered Race, 2023							
0	0	0	0	0	0	0	0				0	0	0	0	0	0	0	0
<1	1-9	10-19	20-29	30-39	40-49	50-59	60+				Asian or Pacific Islander	Black	White	Hispanic	Native American	Other	Unknown	Two or More Races
								Case Count	%	Annual Rate								
								Female	0	0.0								
								Male	0	0.0								
								Unknown	0	0.0								

Psittacosis

Psittacosis is a disease caused by the bacteria *Chlamydia psittaci*. It can be transmitted when a person inhales organisms that have been aerosolized from dried feces or respiratory tract secretions from infected birds. Infected birds might not show signs of illness. Clinical symptoms in people usually consist of fever, headache, weakness, muscle aches, chills, and nonproductive cough. The severity of the disease ranges from a mild, non-specific influenza-like illness to a systemic illness with severe pneumonia. People who own or work with birds should follow precautions when handling and cleaning birds and cages.




No cases of psittacosis were reported in Virginia during 2023.


Age Group, 2023									Sex, 2023				Tiered Race, 2023							
													00000000							
													Asian or Pacific IslanderBlackWhiteHispanicNative AmericanOtherUnknownTwo or More Races							
									Case Count%				Annual Rate							
0									Female				00.0							
1-9									Male				00.0							
10-19									Unknown				00.0							
20-29																				
30-39																				
40-49																				
50-59																				
60+																				

Q fever

Q fever is caused by the bacteria *Coxiella burnetii*. These bacteria naturally infect some animals such as sheep, goats and cattle. People get infected by breathing in dust that has been contaminated by infected animal feces, urine, milk, and birth products that contain *Coxiella burnetii*. Most commonly, people are exposed to this organism via inhalation of infectious aerosols directly from birth fluids of infected animals or via inhalation of dust contaminated with dried birth fluids or tissues. Approximately 50% of people exposed to Q fever will develop the acute form of illness, which is characterized by high fever, severe headache, fatigue, chills, and muscles aches. Chronic Q fever is a severe disease developing in less than 5% of patients exposed. Infection of heart valves is a common form of chronic disease. Preventive measures include appropriate disposal of potentially infectious tissues and proper hygiene when handling animal birth material.




Three cases of Q fever were reported in Virginia in 2023. Two of these cases were acute Q fever and one was a chronic case. One acute case reported contact with sheep, goats, and cattle and the chronic case had previously been diagnosed with acute Q Fever. Virginia typically reports fewer than five Q fever cases each year.




Rabies, human

Rabies is a rhabdovirus of the genus *Lyssavirus*. Rabies infections are most commonly diagnosed in wildlife, especially in certain wild animals such as raccoons, skunks, and foxes. Wildlife is the reservoir of rabies in the United States and any domestic animal cases are a result of spillover from infections in wildlife. Transmission occurs most commonly through the bite of an infected animal but may be transmitted through any method by which virus-infected saliva or central nervous system tissue enters the body. Important prevention methods include vaccinating domestic animals, using animal control to remove stray animals, avoiding handling wildlife and prompt attention to exposures, including post-exposure vaccinations when necessary.




In 2023, 381 animals in Virginia were laboratory-confirmed with rabies, 91% of which were confirmed in wildlife species. Virginia has reported an average of 375 confirmed animal cases of rabies per year since 2014. No human cases of rabies were reported from Virginia in 2023.




Rubella, including congenital rubella syndrome

Rubella is a mild febrile rash illness caused by rubella virus. The virus can be found in nose and throat secretions, such as saliva or nasal mucus, and is spread through sneezing or coughing. Symptoms can include a low-grade fever, sore throat, and a rash that starts on the face and spreads to the rest of the body. In pregnant women, rubella infection can lead to congenital rubella syndrome (CRS) causing miscarriage or serious birth defects in a developing baby. Cases of rubella in the United States decreased dramatically following the initiation of the rubella vaccination program in 1969. Rubella and CRS were declared eliminated from the U.S. in 2004. Rarely, cases of rubella are reported in individuals infected during international travel.




In the last 10 years, no cases of rubella have been reported in Virginia.




Salmonellosis

Salmonellosis is an infection caused by the bacteria *Salmonella*. It is commonly found in the feces of animals raised for food or kept as pets and infected people. The feces can then contaminate raw meats, chicken, eggs, unpasteurized milk and cheese products and other foods, and people can get sick from eating these contaminated foods. Infected persons can spread the bacteria if they do not wash their hands well after using the bathroom or after handling animals and then touching something that other people put in their mouth. Symptoms of salmonellosis include diarrhea, abdominal cramps, headache, fever, and sometimes vomiting. Spread of salmonellosis can be reduced through proper hand washing before and after preparing foods, after using the bathroom or changing diapers, and after handling animals, and ensuring poultry and meats are cooked to appropriate temperatures.




In 2023, 1,260 cases of salmonellosis were reported in Virginia, with a statewide incidence rate of 14.5 per 100,000 population. Fifty-five percent of cases reported their sex as female.




Shiga toxin-producing *Escherichia coli* infection (STEC)

Escherichia coli (*E. Coli*) are bacteria that normally live in the intestines of humans and animals. Most strains do not cause illness; however, strains that produce toxins, such as Shiga toxin-producing *Escherichia coli* (STEC), can lead to diarrheal illness. People and animals infected with STEC shed the bacteria in their feces (stool) which can contaminate surfaces, food, or water. Most people become infected by touching contaminated surfaces and then putting their hands in their mouths, being in contact with farm animals, or by eating or drinking contaminated food. STEC has been associated with people eating contaminated foods, such as undercooked ground beef, raw produce (e.g., leafy greens), raw dough or batter, or drinking unpasteurized (raw) milk or juice. Transmission of this illness can be reduced through proper hand washing, ensuring beef products are thoroughly cooked, and consuming properly pasteurized dairy and juice products.




In 2023, 484 cases of STEC were reported in Virginia, with a statewide incidence rate of 5.6 per 100,000 population. Children aged 1-9 years represent 21% of all cases reported with a statewide incidence rate of 11.1 per 100,000 population.




Shigellosis

Shigellosis is an infection of the intestines caused by a bacterium called *Shigella*. It is easily spread from person to person and sometimes results in outbreaks. People infected with *Shigella* shed the bacteria in their feces (stool). The feces can then contaminate surfaces, food, or water. People can become infected by touching contaminated surfaces, getting the bacteria on their hands and then putting their hands in their mouths, or by eating contaminated food or drinking contaminated water. Symptoms of shigellosis can include mild or severe diarrhea, often with fever and traces of blood in the stool. Transmission of shigellosis can be prevented by careful handwashing with soap and water after using the bathroom or changing diapers, and before or after preparing food.




In 2023, 412 cases of shigellosis were reported in Virginia, with a statewide incidence rate of 4.7 per 100,000 population. Fifty-six percent of cases reported their sex as male.



Smallpox

Smallpox is a serious, contagious, and deadly disease caused by the variola virus. The virus spreads from person to person through direct contact with respiratory droplets from a cough or sneeze, secretions, or skin lesions of an infected person. People with smallpox report a fever and a distinctive skin rash. The mortality rate is approximately 30%, and between 65-80% of survivors are marked with deep pitted scars (pockmarks), most prominent on the face.



Due to the success of smallpox vaccination campaigns, the last natural outbreak of smallpox in the United States occurred in 1949. In 1980, the World Health Assembly declared smallpox eradicated (eliminated), and no cases of naturally occurring smallpox have happened since.

Age Group, 2023

0	0	0	0	0	0	0	0
<1	1-9	10-19	20-29	30-39	40-49	50-59	60+

Sex, 2023


	Case Count	%	Annual Rate
Female	0		0.0
Male	0		0.0
Unknown	0		0.0

Tiered Race, 2023


0	0	0	0	0	0	0	0
Asian or Pacific Islander	Black	White	Hispanic	Native American	Other	Unknown	Two or More Races

Spotted Fever Rickettsiosis (including RMSF)

Spotted fever rickettsiosis (SFR) is a tick-borne illness that may be caused by several different bacteria in the genus *Rickettsia*, including *Rickettsia rickettsii*, the cause of Rocky Mountain spotted fever (RMSF), and *Rickettsia parkeri*, the cause of Tidewater spotted fever. In Virginia, RMSF, the most serious SFR, may be transmitted by either the brown dog tick or the American dog tick. *Rickettsia parkeri* is transmitted primarily by the Gulf Coast tick. RMSF-positive laboratory results may also be caused by other tick-borne species of *Rickettsia* due to cross-reactivity. RMSF is a severe illness with symptoms including fever, rash, headache, and organ failure. If untreated, cases have up to a 25% mortality rate. Tidewater spotted fever is less severe, and cases often present with an eschar (dry, dead wound tissue) at the site of tick bite. The national case definition was updated in 2020.




In 2023, 92 cases of SFR were reported in Virginia, an increase from the 69 cases reported in 2022. Forty-five percent (45%) of these 92 cases occurred among patients aged 60 and older. There was one reported fatality. While a decline in SFR cases was first noticed in 2020, likely influenced by the COVID-19 pandemic, the number of reported SFR cases has since trended upward.




Streptococcal disease, Group A, invasive or toxic shock

Group A streptococcal (GAS) infections, caused by *Streptococcus pyogenes*, can range from minor illnesses to very serious and deadly conditions. Severe and invasive GAS infections occur when bacteria spread to sterile parts of the body, such as the blood or deep muscle tissue. People typically spread the bacteria through respiratory droplets or direct contact. There are certain groups that are at increased risk of serious or invasive GAS infections, including older adults, people who inject drugs, those with immunocompromising conditions, people receiving wound care, and residents of long-term care facilities. Outbreaks of GAS occur periodically in long-term care facilities, such as nursing homes. Invasive group A streptococcal disease (iGAS) and streptococcal toxic shock syndrome (STSS) are reportable conditions in Virginia. STSS can progress rapidly to cause low blood pressure, multiple organ failure, and even death. Prevention of GAS infections include good hand washing, limiting exposure and spread of the bacteria, use of preventive antibiotics in high-risk individuals, and proper wound care.




Rates of invasive group A streptococcal disease in Virginia have been increasing since 2021. Nationally, preliminary CDC 2023 data indicate invasive GAS infections reached a 20-year high. In Virginia, 661 cases were reported to VDH. Invasive GAS cases had an overall incident rate of 7.6 cases per 100,000 population. Persons aged 60 years and older represented 46% of all cases in 2023 at a rate of 15.1 cases per 100,000 population.




Streptococcus pneumoniae, invasive (age less than 5 years)

Invasive pneumococcal disease (IPD) is caused by the bacteria *Streptococcus pneumoniae* and is spread from person to person through respiratory droplets, such as those produced by coughing or sneezing. The bacteria can cause infections in many parts of the body, including the blood, lungs (pneumonia), sinuses, and lining of the brain and spinal cord (meningitis). Symptoms depend on the part of the body affected, and can include fever, cough, shortness of breath, chest pain, stiff neck, joint pain, and chills. Unvaccinated children, the elderly, and people with weakened immune systems are at highest risk for invasive infection. In Virginia, IPD cases are only reported in children under 5 years of age.




Virginia reported 30 IPD cases in 2023. This represents 6.1 cases per 100,000 Virginians less than 5 years of age.




Syphilis, early

Syphilis is a systemic disease caused by the bacterium *Treponema pallidum*. It can be transmitted through vaginal, oral, or anal sex. It can also be prenatally transmitted from a mother to their unborn child. Syphilis is called the “great imitator” as its many possible symptoms can mimic other diseases. Early symptoms may appear 21 days after exposure (range 10-90 days). If left untreated, syphilis can progress to more severe disease. Syphilis infections disproportionately affect men who have sex with men (MSM), non-Hispanic black persons, and those 20-39 years of age. This report includes all cases of early syphilis (primary, secondary, and early non-primary, non-secondary stages) reported in Virginia. Transmission of this disease can be mitigated by condom use, routine screening for sexually active persons, and ensuring adequate treatment for both infected individuals and their sex partners.




In 2023, 1,764 early syphilis cases were reported, which is a 2% increase in incidence rate from 2022 (19.9 to 20.3 per 100,000). Rates continue to be highest among males (33.6 per 100,000) and cases among men account for 82% of all reported cases, yet the rate from 2022 to 2023 increased 23% in females compared to a 1% decrease in males. Rates of early syphilis also continue to be highest among Black persons (57.9 per 100,000) and persons aged 30 to 39 years (51.6 per 100,000).



Tetanus

Tetanus, commonly known as lockjaw, is a disease caused by *Clostridium tetani* bacteria that affects the nervous system. The bacteria can enter the body through a break in the skin, primarily a wound, and produce a toxin that causes painful muscle contractions. Signs and symptoms can include jaw cramping, muscle spasms, trouble swallowing, seizures, headache, and fever. The risk of death from tetanus is highest among people 60 years of age or older. Since 1947, reported cases of tetanus have declined more than 95% in the United States due to widespread use of the tetanus vaccine. To this day, cases of tetanus remain rare in the United States.




No cases of tetanus were reported in Virginia in 2023.

Age Group, 2023									Sex, 2023				Tiered Race, 2023							
													0 0 0 0 0 0 0 0							
													</							

Trichinosis (Trichinellosis)

Trichinellosis, also known as trichinosis, is caused by eating raw or undercooked meat that contains the early, immature form (larvae) of a worm called *Trichinella*. It occurs worldwide, most often in wild animals, but it is not common in the United States. People who eat raw or undercooked meat from animals infected with the *Trichinella* worm are at higher risk for developing trichinellosis. Meats of concern include bear, pork, wild feline (such as cougar), fox, dog, wolf, horse, seal, and walrus. The disease is not spread from person to person. Initial symptoms of trichinellosis include abdominal discomfort, nausea, vomiting, diarrhea, fatigue, and fever. Aching joints, muscle pain, swelling of the face and eyes, sweating, chills, headaches, cough, itchy skin, and sometimes constipation can occur later.



No cases of trichinosis were reported in 2023.


Age Group, 2023							
<1	1-9	10-19	20-29	30-39	40-49	50-59	60+
0	0	0	0	0	0	0	0

Sex, 2023			
	Case Count	%	Annual Rate
Female	0		0.0
Male	0		0.0
Unknown	0		0.0


Tiered Race, 2023						
0	0	0	0	0	0	0
Asian or Pacific Islander	Black	White	Hispanic	Native American	Other	Two or More Races

Tuberculosis

Tuberculosis (TB) is a disease caused by the bacterium *Mycobacterium tuberculosis*. TB most often affects the lungs, but can occur anywhere in the body. There are two TB-related conditions: TB disease and latent TB infection (LTBI). People with TB disease usually feel sick, have symptoms, and can spread TB to others, depending on the site of disease. Over 80% of people with TB in Virginia are born outside of the United States, and international humanitarian crises and global migration impacts TB burden. Prompt initiation of treatment and contact investigations reduces transmission.




In 2023, Virginia documented a six percent increase in people diagnosed with TB disease compared to 2022. This increase was consistent with increases seen nationally with a return to pre-pandemic trends. Virginia's TB rate stayed below the national average, but increased slightly from prior years.




Tularemia

Tularemia is a disease caused by the bacteria *Francisella tularensis*. Tularemia can be transmitted in a number of ways, including tick bites, handling an infected animal, consuming contaminated water, and laboratory exposure. Symptoms can vary depending on the mode of transmission, but usually include sudden onset of high fever, chills, fatigue, general body aches, headache, and nausea. Preventive measures include minimizing the risk of tick bites by the use of both appropriate dress and insect repellents when recreating or working in tick habitats and avoiding the consumption of untreated water. Impervious protective gloves should be used when skinning rabbits and other wild game. Utensils used for preparing meat from game should not be used to prepare other food items. Undercooked meat should not be consumed. Mowing over dead animals should be avoided to lower the risk of aerosolizing infectious particles.




Four cases of tularemia were reported in 2023. All four cases were hospitalized, and three cases appeared to be tick bite related. Typically, Virginia reports fewer than 5 cases per year.




Typhoid infection

Typhoid infection is a serious disease caused by the bacterium *Salmonella* Typhi. Most cases diagnosed in the U.S. are acquired during travel to other countries in Asia, Africa, and South and Central America, where the disease is common. It can be transmitted by eating or drinking food or water that has been contaminated with feces or urine of people with the disease or by direct contact with a person who has the disease. Typhoid infection can cause high fever, headaches, weakness, loss of appetite, diarrhea, or constipation, and some people get “rose spots” on the trunk of the body. Typhoid infection can be prevented by proper hand washing after using a toilet or changing a diaper, and before preparing or eating food. A vaccine is available that provides some protection for persons traveling to areas where the disease is common. Even if vaccinated, persons traveling to these areas still need to be careful about what food and water they consume.




In 2023, 15 cases of typhoid infection were reported in Virginia, with a statewide incidence rate of 0.2 per 100,000 population. Children aged 1-9 years represented 40% of cases reported. Sixty-six percent of cases reported their sex as female.




Vaccinia adverse event following vaccination

The smallpox vaccine is made from the live vaccinia virus, which is similar to the variola virus that causes smallpox but less harmful. Routine smallpox vaccination stopped after the disease was eradicated. Vaccination is still recommended for people who are at high risk of occupational exposure, such as laboratory staff who handle smallpox virus and other orthopoxviruses. Most vaccinated people will have a mild reaction, such as a sore arm, fever, swollen glands, and body aches. More severe reactions can occur, including dermatologic complications or cardiac adverse events, and must be reported to public health. Also, vaccinia disease or adverse events can occur in close contacts exposed to the fluid or crust material from the lesion of someone who was recently vaccinated. People who were recently vaccinated must take precautions to prevent vaccinia virus from spreading.




The last reported case of vaccinia disease or adverse event was reported in 2018. No cases were reported in 2023.




Vancomycin-intermediate *Staphylococcus aureus* (VISA)

Staphylococcus aureus is a bacterium commonly found on the skin and in the nose. Spread occurs among people having close physical contact with infected patients or contaminated material, such as bandages. Early detection and implementation of infection prevention and control strategies are necessary to prevent the spread. Sometimes these bacteria can get into the bloodstream and cause serious infections which can be fatal. Vancomycin-intermediate *Staphylococcus aureus* is a type of staph infection that has intermediate resistance to a drug called vancomycin. Vancomycin is a common treatment option for staph infections. Persons who develop this type of staph infection may have underlying health conditions, tubes going into their bodies, previous infections with methicillin-resistant *Staphylococcus aureus* (MRSA), and recent exposure to vancomycin and other antimicrobial agents.




VISA is rare in Virginia with no cases reported in 2023.




Vancomycin-resistant *Staphylococcus aureus* (VRSA)

Staphylococcus aureus is a bacterium commonly found on the skin and in the nose. Spread occurs among people having close physical contact with infected patients or contaminated material, such as bandages. Early detection and implementation of infection prevention and control strategies are necessary to prevent the spread. Sometimes these bacteria can get into the bloodstream and cause serious infections which can be fatal. Vancomycin-resistant *Staphylococcus aureus* is a type of staph infection that has resistance to a drug called vancomycin. Vancomycin is a common treatment option for staph infections. Persons who develop this type of staph infection may have underlying health conditions, tubes going into their bodies, previous infections with methicillin-resistant *Staphylococcus aureus* (MRSA), and recent exposure to vancomycin and other antimicrobial agents.




VRSA is extremely rare in the United States and Virginia has not identified a case of VRSA.




Vibriosis, non-cholera

Vibriosis is a potentially serious illness caused by a group of bacteria called *Vibrio*. Anyone can be infected with *Vibrio*, but people with weakened immune systems and conditions that damage the liver are more likely to experience severe illness. *Vibrio* bacteria can cause different types of infection including gastrointestinal, wound, and blood. Most people become infected through eating raw or undercooked seafood and shellfish or when the *Vibrio* bacteria enter the body through a break in the skin while a person is in salt or brackish waters. Vibriosis is most common during summer months when the waters are warmer.




In 2023, 117 cases of vibriosis were reported in Virginia, with a statewide incidence rate of 1.3 per 100,000 population. Adults 60 years of age and older represented 37% of cases reported. Fifty-five percent of cases reported their sex as female.




Viral hemorrhagic fever

Viral hemorrhagic fevers (VHFs) are a group of diseases that are caused by several distinct families of viruses. The term “viral hemorrhagic fever” refers to a condition that affects many organ systems of the body, damages the overall cardiovascular system, and reduces the body’s ability to function on its own. Symptoms vary but often include bleeding or hemorrhaging. Illness severity is wide ranging, from relatively mild illness to severe, life-threatening disease. Most VHFs have no known cure or vaccine. VHF viruses can spread to people when they come in contact with infected animals or insects. For many VHFs, person-to-person transmission can then continue. Outbreaks of VHFs occur sporadically and irregularly, and their occurrence can be difficult to predict. Prevention is more difficult when the animal host is unknown or challenging to control (such as rodents or ticks).




No cases of viral hemorrhagic fever, of any kind, were reported in Virginia in 2023.




Yellow fever

Yellow fever is caused by a virus that is primarily found in tropical and subtropical areas of Africa and South America. The virus is transmitted through the bites of several species of infected *Aedes* mosquitoes, most notably the yellow fever mosquito (*Aedes aegypti*), which breeds in containers of water occurring around human habitats. Yellow fever mosquitoes occur in Virginia but have become rare after being displaced from their container breeding habitats by the arrival of the closely related Asian tiger mosquito (*Aedes albopictus*) in 1992. Yellow fever is a very rare cause of illness in U.S. travelers. Illness ranges from a fever with aches and pains to severe liver disease with bleeding and yellowing skin (jaundice). Vaccination against the yellow fever virus before traveling to endemic regions, and avoidance of mosquito bites while traveling in these regions, can help prevent infections.



Yersiniosis

Yersiniosis is a gastrointestinal disease caused by the bacteria *Yersinia*. The bacteria can be found in live animals, particularly pigs. It can be transmitted by eating contaminated food, especially raw or undercooked pork products. Surfaces can become contaminated with *Yersinia* when preparing raw pork. If another food item or hands touch those surfaces and enter someone's mouth, the person can become infected with the bacteria. Yersiniosis can cause fever, abdominal pain, and diarrhea. Transmission of this condition can be reduced through proper hand washing and ensuring pork and pork products are cooked to appropriate temperatures.



In 2023, 192 yersiniosis cases were reported in Virginia with a statewide incidence rate of 2.2 per 100,000 population. This represents a 32% increase over the 145 cases reported in 2022. Adults 60 years of age and older represented 42% of yersiniosis cases and 63% reported their sex as female.

