

# Test Your Knowledge

## 1. TRUE OR FALSE

INSULIN PENS CAN BE USED FOR MULTIPLE PATIENTS/RESIDENTS.

## 2. TRUE OR FALSE

SINGLE-USE, AUTO-DISABLING FINGERSTICK SHOULD ALWAYS BE USED WHEN ASSISTED BLOOD GLUCOSE MONITORING IS PERFORMED

## 3. TRUE OR FALSE

HAND HYGIENE IS PERFORMED BEFORE AND AFTER PERFORMING A FINGERSTICK PROCEDURE TO OBTAIN A SAMPLE OF BLOOD AND WHEN USING THE GLUCOMETER (OR OTHER POC DEVICE).

4. IF A BLOOD GLUCOSE METER (OR GLUCOMETER) IS USED FOR MORE THAN ONE PATIENT WHAT MUST BE DONE?

- A. BGM properly cleaned with soap and water
- B. BGM wiped down with a sani wipe
- C. Follow manufacturer's instructions to clean and disinfect the BGM with appropriate disinfectant and dwell time after each use
- D. Nothing needs to be done

5. WHEN SHOULD EMPLOYEES PROVIDING AMBG RECEIVE TRAINING ON PROPER USE OF DEVICES (E.G., LANCETS, INSULIN PENS, AND BLOOD GLUCOSE MONITORS)?

- A. Upon Hire
- B. Annually
- C. New Device is employed
- D. All of the above

6. WHAT IS THE PROPER WAY TO LABEL A RESIDENT'S INSULIN PEN?

- A. With patient's name and home address
- B. With the patient's date of birth
- C. With the expiration date
- D. With the patient's name and other identifiers

7. WHICH OF THE FOLLOWING IS AN UNSAFE INJECTION PRACTICE?

- A. Using the same syringe to administer medication to more than one patient
- B. Using medications packaged as single-dose or single-use for one patient
- C. Using aseptic technique when preparing and administering injections
- D. Performing hand hygiene prior to and after preparation and administration of medication