Network Systems Science & Advanced Computing

Biocomplexity Institute & Initiative

University of Virginia

Estimation of COVID-19 Impact in Virginia

July 8th, 2020

(data current to July 7th)

Biocomplexity Institute Technical report: TR 2020-084

UNIVERSITY of VIRGINIA

BIOCOMPLEXITY INSTITUTE

biocomplexity.virginia.edu

Who We Are

- Biocomplexity Institute at the University of Virginia
  - Using big data and simulations to understand massively interactive systems and solve societal problems
- Over 20 years of crafting and analyzing infectious disease models
  - Pandemic response for Influenza, Ebola, Zika, and others

Points of Contact

Bryan Lewis
brylew@virginia.edu

Srini Venkatramanan
srini@virginia.edu

Madhav Marathe
marathe@virginia.edu

Chris Barrett
ChrisBarrett@virginia.edu

Biocomplexity COVID-19 Response Team

Aniruddha Adiga, Abhijin Adiga, Hannah Baek, Chris Barrett, Golda Barrow, Richard Beckman, Parantapa Bhattacharya, Andrei Bura, Jiangzhuo Chen, Clark Cucinell, Allan Dickerman, Stephen Eubank, Arindam Fadikar, Joshua Goldstein, Stefan Hoops, Sallie Keller, Ron Kenyon, Brian Klahn, Gizem Korkmaz, Vicki Lancaster, Bryan Lewis, Dustin Machi, Chunhong Mao, Achla Marathe, Madhav Marathe, Fanchao Meng, Henning Mortveit, Mark Orr, Przemyslaw Porebski, SS Ravi, Erin Raymond, Jose Bayoan Santiago Calderon, James Schlitt, Aaron Schroeder, Stephanie Shipp, Samarth Swarup, Alex Telionis, Srinivasan Venkatramanan, Anil Vullikanti, James Walke, Amanda Wilson, Dawen Xie

UNIVERSITY of VIRGINIA

BIOCOMPLEXITY INSTITUTE

Overview

- **Goal**: Understand impact of COVID-19 mitigations in Virginia
- **Approach:**
  - Calibrate explanatory mechanistic model to observed cases
  - Project infections through the end of summer
  - Consider a range of possible mitigation effects in "what-if" scenarios
- **Outcomes:**
  - Ill, Confirmed, Hospitalized, ICU, Ventilated, Death
  - Geographic spread over time, case counts, healthcare burdens

Key Takeaways

Projecting future cases precisely is impossible and unnecessary.
Even without perfect projections, we can confidently draw conclusions:

- **Early indicators that some VDH health districts are seeing increased activity. With other regions of the country experiencing resurgence, it is crucial to maintain control.**
- Recent model updates:
  - Integrating hospital occupancy data for model validation
  - Updated potential resurgence scenario
  - Identified “Best fitting” scenarios per health district based on recent trends and added a combined state level “Best Fit” scenario representing this combination
  - Updated additional analyses to inform restructuring of model for next phase of epidemic
- Other states showing rapid rise following relaxation of social distancing with limited control measures.
- The situation is changing rapidly. Models will be updated regularly.

Model Configuration and Data Analysis

Simulation Engine – PatchSim

- Metapopulation model
  - Represents each population and its interactions as a single patch
  - 133 patches for Virginia counties and independent cities
- Extended SEIR disease representation
  - Includes asymptomatic infections and treatments
- Mitigations affect both disease dynamics and population interactions
- Runs fast on high-performance computers
  - Ideal for calibration and optimization





Venkatramanan, Srinivasan, et al. "Optimizing spatial allocation of seasonal influenza vaccine under temporal constraints." *PLoS Computational Biology* 15.9 (2019): e1007111.

Model Configuration

- **Transmission:** Parameters are calibrated to the observed case counts
  - **Reproductive number:** 2.1 - 2.3
  - **Infectious period** (time of infectiousness before full isolation): 3.3 to 5 days
- **Initial infections:** Start infections from confirmed cases by county
  - Timing and location based on onset of illness from VDH data
  - Assume 15% detection rate, so one confirmed case becomes ~7 initial infections
- **Mitigations:** Intensity of social distancing rebound and control sustaining mitigations into the future are unknowable, thus explored through 5 scenarios

Mitigation Scenarios: Rebound Intensity x Detection

Pause from Social Distancing: Began on March 15th

- Lifted on May 15th (61 days), with two-week delay (75 days) for select counties*
- **Intensity**: Social distancing pauses and significantly reduces case growth, this level varies by VDH Health District and is fit through an analysis of growth rate during the Pause

Intensity of Rebound:

- **Steady:** Intensity of effective mixing remains steady from Pause as infection control practices moderate increased interactions
- **Light:** Effective mixing returns to 1/6th of pre-pandemic levels
- **Full Rebound**: Interactions return completely (100%) to pre-pandemic levels, as a reference

Tracing and Isolation: Increased Testing Capacity coupled with infection control measures can limit the period of infectiousness without isolation

- **Better Detection**: Observed relative reductions in days from onset to diagnosis applied to infectious period from (30% and then 45%) and remain stable into future for projections

* Select counties as mentioned by recent releases from Governor Northam's office
https://www.governor.virginia.gov/newsroom/all-releases/2020/may/headline-856741-en.html
https://www.governor.virginia.gov/newsroom/all-releases/2020/may/headline-856796-en.html

Five Mitigation Scenarios

Scenario	Rebound Intensity	Better Detection	Name	Description
1	Light	No	Light	Light Rebound, Detection same
2	Steady	No	Steady	Steady Interactions, Detection same
3	Light	Yes	Light – BetterDetection	Light Rebound, Detection improved
4	Steady	Yes	Steady – BetterDetection	Steady Interactions, Detection improved
5	Full	No	Full Rebound	Return to No mitigation

Potential Future Surge Scenario

Resurgence: Several other states are experiencing a resurgence in cases

- Analysis of of 23 states with some level or resurgence shows average time from most complete relaxation of social distancing and resurgence in cases was 28 days
- Not certain if or when this may occur in Virginia, this projection uses 28 days

Intensity of Surge:

- Difficult to predict
- Using Strong Rebound, 1/3 back to pre-pandemic level for transmission
- "Better Detection" scenarios maintain recent observed increases

Surge Projection: 4 weeks from July 1st, increase in transmissibility and recent increase in delay of case detection remains steady



UNIVERSITY of VIRGINIA
BIOCOMPLEXITY INSTITUTE

Full Model Parameters

	Parameter	Values	Description
Transmission	Transmissibility (R_0)[1]	2.2 [2.1 – 2.3]	Reproductive number
	Incubation period[1]	5 days	Time from infection to infectious
	Infectious period[1]	3.3 - 5 days	Duration of infectiousness
	Infection detection rate[3]	15%	1 confirmed case becomes ~7 initial infections
	Percent asymptomatic[1]	50%	Infected individuals that don't exhibit symptoms
Resources	Onset to hospitalization[1]	5 days	Time from symptoms to hospitalization
	Hospitalization to ventilation[1]	3 days	Time from hospitalization to ventilation
	Duration hospitalized	8 days	Time spent in the hospital [4]
	Duration ventilated[2]	14 days	Time spent on a ventilator
	Percent hospitalized[1]	5.5% (~20% of confirmed)	Symptomatic individuals becoming hospitalized
	Percent in ICU[1]	20%	Hospitalized patients that require ICU
	Percent ventilated[1]	70%	ICU patients requiring ventilation
	Percent Fatality	1.35%	Symptomatic individuals who die

1 CDC COVID-19 Modeling Team. "Best Guess" scenario. Planning Parameters for COVID-19 Outbreak Scenarios. Version: 2020-03-31.
2 Up-to-date. COVID-19 Critical Care Issues. https://www.uptodate.com/contents/coronavirus-disease-2019-covid-19-critical-care-issues?source=related_link (Accessed 13APRIL2020)
3 Li et al., *Science* 16 Mar 2020:eabb3221 https://science.sciencemag.org/content/early/2020/03/24/science.abb3221 (Accessed 13APRIL2020)
4 Personal communications, UVA Health and Sentara (~500 VA based COVID patients)

Calibration Approach

- **Data:**
  - County level case counts by date of onset (from VDH)
  - Confirmed cases for model fitting
- **Model:** PatchSim initialized with disease parameter ranges from literature
- **Calibration:** fit model to observed data
  - Search transmissibility and duration of infectiousness
  - Markov Chain Monte Carlo (MCMC) particle filtering finds best fits while capturing uncertainty in parameter estimates
- **Spatial Adjustments:** VDH districts grouped to 3 tiers of growth during the Pause, with similarly scaled reductions then applied to the groups of districts
- **Project:** future cases and outcomes using the trained particles



Accessed 11pm July 7, 2020
https://www.vdh.virginia.gov/coronavirus/

Spatial Adjustments at District Level

Case Rate (per 100K) by VDH District

- Regions arranged to rough position in Commonwealth and colored by VDH Health Region
- Considerable variation across districts
- Some consistent behaviors during mid-April to mid-May during the Pause period
- Smoothed (Savitzky-Golay filter)



Spatial Adjustments at District Level

Adjustments based on Growth during Pause

- Group districts by their mean growth from mid-April to mid-May (using model based R_{eff})
- Assign reductions during Pause, and beyond, to members of these groups
- Low reduction = 40%
- Moderate reduction = 45% (previous level)
- High reduction = 55%



Impact of Interventions

Estimating Effects of Social Distancing

Mobility data shows pause mid-March, slow rebound starting in May

Google Mobility data shows continued slow rebound
(as of June 14th)
https://www.google.com/covid19/mobility/

- Regional levels of Stay at home vs. Workplace
- 35% reduction of those staying at home, very slow and stable reductions
- Other activities show vaster increases with grocery / retail nearly back to baseline
- Parks and recreation show significant increase





Weekly growth rate by date of onset

- Week before March 15 = 0.3
- Week after March 15 = -0.03 to 0.04

Crude reproductive number estimates

- 2.2 before March 15th
- 0.81 to 1.10 after March 15th



Estimating Changes in Case Detection

VDH data show changes in time from Symptom Onset to Diagnosis

Days to Diagnosis dropped but rebounding

- Mid March to Late April = 7.8 days
- Late April to Mid May = 5.7 days (~25% lower)
- Mid May to early June = 4.7 days (~40% lower)
- Early June to late June = 5.1 days (~35% lower)

Test positivity vs. Onset to Diagnosis



Recent flattening / slight rise in test positivity

Testing Encounters have steadied and increased



Recent flattening then slight rise in total tests

Accessed 9pm July 7th, 2020
https://www.vdh.virginia.gov/coronavirus/



Estimating Changes in Case Detection* – by district



*up to the end of June when data is stable

UNIVERSITY of VIRGINIA
BIOCOMPLEXITY INSTITUTE

Estimating Daily Reproductive Number





Statewide and most regions follow similar pattern

- Spike, followed by a decline, to plateau, with recent upswing
- This week: overall decline, some regions up

Methodology

- Wallinga-Teunis method as implemented in EpiEstim[1] R package
- Based on Date of Onset of Symptoms
- Uses serial interval to estimate R_e over time: 6 days (2 day std dev)

Recent Estimates subject to revision as more data comes in

- Date of onset unstable in last 7-14 days

June 27th Estimates

Region	Current R_e	Diff Last Week
State-wide	1.124	0.187
Central	0.911	0.089
Eastern	1.399	0.216
Far SW	1.935	1.223
Near SW	1.228	-0.422
Northern	1.103	0.313
Northwest	0.993	-0.156



1. Anne Cori, Neil M. Ferguson, Christophe Fraser, Simon Cauchemez. A New Framework and Software to Estimate Time-Varying Reproduction Numbers During Epidemics. American Journal of Epidemiology, Volume 178, Issue 9, 1 November 2013, Pages 1505–1512, https://doi.org/10.1093/aje/kwt133

Impact of Race / Ethnicity & Outbreaks

Confirmed Case Rate



Death Rate





Different Races and Ethnicities disproportionally affected

- Hispanic population bears large burden of disease compared to population size

Outbreak Events are hard to predict and affect model fits

- Eastern Shore has 60% of cases from 10 outbreaks
- Fairfax most outbreaks but relatively low proportion

Other State Comparisons











Several States experiencing large surges in cases

- Of the 23 with significant upticks we calculated an average of 28 days from policy guidance to more fully relax social distancing and the resurgence of cases

Re Estimates for VA and neighbors show upward trend

- Virginia above 1, joining NC and TN
- DC and MD still below 1 but trending up

Early signs of resurgence: Plateauing or increase in test positivity and R above 1 for several weeks

Estimated Re* for surging States and Neighbors



* Based on confirmed cases per day

UNIVERSITY of VIRGINIA

BIOCOMPLEXITY INSTITUTE

Model Results

Outcome Projections

Confirmed cases



Hospital occupancy



Deaths



Cumulative Confirmed cases



UNIVERSITY of VIRGINIA
BIOCOMPLEXITY INSTITUTE

Outcome Projections – with Surge



Weekly New Confirmed Cases*

Week Ending	Best Fit	Best Fit w/ Surge
6/28/20	4,045	4,045
7/5/20	3,621	3,621
7/12/20	3,291	3,291
7/19/20	3,375	3,375
7/26/20	3,492	3,492
8/2/20	3,570	3,676
8/9/20	3,628	4,836
8/16/20	3,692	6,001
8/23/20	3,740	7,438
8/30/20	3,760	9,192
9/6/20	3,758	11,208

*Numbers are medians of projections

UNIVERSITY of VIRGINIA
BIOCOMPLEXITY INSTITUTE

Hospital Demand and Capacity by Region

Capacities by Region – BestFit- Surge

COVID-19 capacity ranges from 80% (dots) to 120% (dash) of total beds



- Based on current best fits with potential surge
  - No region exceeds 80% capacity until end of August
  - Multiple regions (Near SW, Far SW, Eastern and Northern) may near their capacity in September
- Next few weeks (until mid-August) are crucial to mitigate/prepare for a surge in cases
- Activity in neighboring states and reopening of schools/universities may make this more likely

* Assumes average length of stay of 8 days

District Level Projections - Daily

Best fitting projections by District

- Projections that best fit recent trends
- Daily confirmed cases by Region (blue solid) with simulation at the region level (black dotted)
- Projection color consistent with other and abbreviated in title



Abbrv	Name	# of Districts (last wk)
LR	Light	2 (1)
S	Steady	12 (9)
LR-BD	Light – BetterDetection	14 (9)
S-BD	Steady – BetterDetection	7 (16)
FR	Full Rebound	0

District Level Projections – Daily with Surge

Best fitting projections by District

- Projections that best fit recent trends
- Daily confirmed cases by Region (blue solid) with simulation at the region level (black dotted)
- Projection color consistent with other and abbreviated in title



Abbrv	Name	# of Districts (last wk)
LR	Light	2 (1)
S	Steady	12 (9)
LR-BD	Light – BetterDetection	14 (9)
S-BD	Steady – BetterDetection	7 (16)
FR	Full Rebound	0

District Level Projections – Cumulative

Best fitting projections by District

- Projections that best fit recent trends
- Daily confirmed cases by Region (blue solid) with simulation at the region level (black dotted)
- Projection color consistent with other and abbreviated in title



Abbrv	Name	# of Districts (last wk)
LR	Light	2 (1)
S	Steady	12 (9)
LR-BD	Light – BetterDetection	14 (9)
S-BD	Steady – BetterDetection	7 (16)
FR	Full Rebound	0

District Level Projections – Cumulative with Surge

Best fitting projections by District

- Projections that best fit recent trends
- Daily confirmed cases by Region (blue solid) with simulation at the region level (black dotted)
- Projection color consistent with other and abbreviated in title



Abbrv	Name	# of Districts (last wk)
LR	Light	2 (1)
S	Steady	12 (9)
LR-BD	Light – BetterDetection	14 (9)
S-BD	Steady – BetterDetection	7 (16)
FR	Full Rebound	0

Key Takeaways

Projecting future cases precisely is impossible and unnecessary.
Even without perfect projections, we can confidently draw conclusions:

- **Early indicators that some VDH health districts are seeing increased activity. With other regions of the country experiencing resurgence, it is crucial to maintain control.**
- Recent model updates:
  - Integrating hospital occupancy data for model validation
  - Updated potential resurgence scenario
  - Identified “Best fitting” scenarios per health district based on recent trends and added a combined state level “Best Fit” scenario representing this combination
  - Updated additional analyses to inform restructuring of model for next phase of epidemic
- Other states showing rapid rise following relaxation of social distancing with limited control measures.
- The situation is changing rapidly. Models will be updated regularly.

References

Venkatramanan, S., et al. "Optimizing spatial allocation of seasonal influenza vaccine under temporal constraints." *PLoS computational biology* 15.9 (2019): e1007111.

Arindam Fadikar, Dave Higdon, Jiangzhuo Chen, Bryan Lewis, Srinivasan Venkatramanan, and Madhav Marathe. Calibrating a stochastic, agent-based model using quantile-based emulation. SIAM/ASA Journal on Uncertainty Quantification, 6(4):1685–1706, 2018.

Adiga, Aniruddha, Srinivasan Venkatramanan, Akhil Peddireddy, et al. "Evaluating the impact of international airline suspensions on COVID-19 direct importation risk." *medRxiv* (2020)

NSSAC. PatchSim: Code for simulating the metapopulation SEIR model. https://github.com/NSSAC/PatchSim (Accessed on 04/10/2020).

Virginia Department of Health. COVID-19 in Virginia. http://www.vdh.virginia.gov/coronavirus/ (Accessed on 04/10/2020)

Biocomplexity Institute. COVID-19 Surveillance Dashboard. https://nssac.bii.virginia.edu/covid-19/dashboard/

Google. COVID-19 community mobility reports. https://www.google.com/covid19/mobility/

Cuebiq: COVID-19 Mobility insights. https://www.cuebiq.com/visitation-insights-covid19/

Biocomplexity page for data and other resources related to COVID-19: https://covid19.biocomplexity.virginia.edu/

Questions?

Points of Contact

Bryan Lewis
brylew@virginia.edu

Srini Venkatramanan
srini@virginia.edu

Madhav Marathe
marathe@virginia.edu

Chris Barrett
ChrisBarrett@virginia.edu

Biocomplexity COVID-19 Response Team

Aniruddha Adiga, Abhijin Adiga, Hannah Baek, Chris Barrett, Golda Barrow, Richard Beckman, Parantapa Bhattacharya, Andrei Bura, Jiangzhuo Chen, Clark Cucinell, Allan Dickerman, Stephen Eubank, Arindam Fadikar, Joshua Goldstein, Stefan Hoops, Sallie Keller, Ron Kenyon, Brian Klahn, Gizem Korkmaz, Vicki Lancaster, Bryan Lewis, Dustin Machi, Chunhong Mao, Achla Marathe, Madhav Marathe, Fanchao Meng, Henning Mortveit, Mark Orr, Przemyslaw Porebski, SS Ravi, Erin Raymond, Jose Bayoan Santiago Calderon, James Schlitt, Aaron Schroeder, Stephanie Shipp, Samarth Swarup, Alex Telionis, Srinivasan Venkatramanan, Anil Vullikanti, James Walke, Amanda Wilson, Dawen Xie

UNIVERSITY of VIRGINIA

BIOCOMPLEXITY INSTITUTE

Supplemental Slides

Recent Parameter Validation

New York State announced sero-prevalence survey results on May 2nd

- 15,000 antibody tests conducted randomly through the state at grocery stores
- **Total Attack Rate**: 12.3%

Estimation of undetected infections

- Total infections in NY = 2.46M, total of 300K confirmed cases
- Confirmed case detection = 12% of infections (close to 15% used in model)

Estimation of hospitalizations from infections

- Total infections in NY = 2.46M, total of 66K hospitalizations
- Hospitalizations = 2.7% of infections (close to 2.25% used in model)

Agent-based Model (ABM)

EpiHiper: Distributed network-based stochastic disease transmission simulations

- Assess the impact on transmission under different conditions
- Assess the impacts of contact tracing



Synthetic Population

- Census derived age and household structure
- Time-Use survey driven activities at appropriate locations



Detailed Disease Course of COVID-19

- Literature based probabilities of outcomes with appropriate delays
- Varying levels of infectiousness
- Hypothetical treatments for future developments

ABM Social Distancing Rebound Study Design

Study of ”Stay Home” policy adherence

- Calibration to current state in epidemic
- Implement “release” of different proportions of people from ”staying at home”



	date	best case	best guess	worst case	R
1	2020-05-10	664	852	1079	0.95
2	2020-05-17	666	841	1093	0.91
3	2020-05-24	617	768	1111	0.80
4	2020-05-31	547	705	1064	NA

Calibration to Current State

- Adjust transmission and adherence to current policies to current observations
- For Virginia, with same seeding approach as PatchSim



Impacts on Reproductive number with release

- After release, spike in transmission driven by additional interactions at work, retail, and other
- At 25% release (70-80% remain compliant)
- Translates to 15% increase in transmission, which represents a 1/6th return to pre-pandemic levels

Medical Resource Demand Dashboard

https://nssac.bii.virginia.edu/covid-19/vmrddash/



UNIVERSITY of VIRGINIA
BIOCOMPLEXITY INSTITUTE