

Common Pediatric Pulmonary Issues

Chris Woleben MD, FAAP
Associate Dean, Student Affairs
VCU School of Medicine
Assistant Professor, Emergency Medicine and Pediatrics

Objectives


- Learn common causes of upper and lower airway disease in the pediatric population
- Learn basic management skills for common pediatric pulmonary problems

Upper Airway Disease

- Extrathoracic structures
 - Pharynx, larynx, trachea
- Stridor
 - Externally audible sound produced by turbulent flow through narrowed airway
 - Signifies partial airway obstruction
 - May be acute or chronic

Remember Physics?

Poiseuille's Law



Acute Stridor

- Febrile
 - Laryngotracheitis (croup)
 - Retropharyngeal abscess
 - Epiglottitis
 - Bacterial tracheitis
- Afebrile
 - Foreign body
 - Caustic or thermal airway injury
 - Angioedema

Croup - Epidemiology

- Usually 6 to 36 months old
- Males > Females (3:2)
- Fall / Winter predilection
- Common causes:
 - Parainfluenza
 - RSV
 - Adenovirus
 - Influenza

Croup - Pathophysiology

- Begins with URI symptoms and fever
- Infection spreads from nasopharynx to larynx and trachea
- Subglottic mucosal swelling and secretions lead to narrowed airway
- Development of barky, “seal-like” cough with inspiratory stridor
- Symptoms worse at night


Croup - Management

- Keep child as calm as possible, usually sitting in parent's lap
- Humidified saline via nebulizer
- Steroids (Dexamethasone 0.6 mg/kg)
 - Oral and IM route both acceptable
- Racemic Epinephrine
 - <10kg: 0.25 mg via nebulizer
 - >10kg: 0.5 mg via nebulizer

Croup – Management

- Must observe for 4 hours after use of racemic epinephrine
- Admit if patient has recurrent stridor or any signs of respiratory distress
- Consider AP/Lateral neck films
 - Steeple sign

Croup – Steeple Sign



Retropharyngeal Abscess

- Etiology
 - Prior pharyngitis, otitis
 - Penetrating wound to posterior pharynx
- Pathophysiology
 - Cellulitis and suppurative adenitis of lymph node in prevertebral fascia

Retropharyngeal Abscess

- Presentation:
 - Fever
 - Difficulty swallowing
 - Drooling
 - Sore throat
 - Changes in voice
 - Stiff neck

Retropharyngeal Abscess



Retropharyngeal Abscess

- Diagnosis
 - Lateral soft tissue neck X-ray
 - Retropharyngeal soft tissue at level of C-3 is greater than 5mm
 - Retropharyngeal soft tissue is more than 40% of the body of C-4 at that level
 - Soft tissue neck CT
 - Better delineate extent of lesion

Retropharyngeal Abscess



Retropharyngeal Abscess

- Management:
 - Assess and secure patent airway
 - Antibiotic coverage
 - Nafcillin and Clindamycin
 - Analgesia
 - ENT consult for operative incision and drainage of abscess


Epiglottitis

- Life threatening emergency!!!
- Clinical presentation:
 - Sudden onset high fever
 - Moderate to severe respiratory distress
 - Stridor
 - Drooling
 - Toxic appearing child
 - Sits leaning forward in a sniffing position with an open mouth

Epiglottitis

- Not seen as frequently today
 - *Haemophilus influenzae* type B vaccine
- Other bacterial causes include staphylococcus and streptococcus
- Diagnosis:
 - Lateral soft tissue neck
 - Epiglottis is rounded and blurred (thumbprint sign)

Epiglottitis



Epiglottitis




Figure 14.38 Epiglottitis

Epiglottitis

- Management:
 - Keep child as calm as possible, preferably in parent's lap
 - Surgical consult to establish definitive airway in operating room
 - Start broad spectrum antibiotic coverage
 - Second or third generation cephalosporins


Bacterial Tracheitis

- Bacterial complication of a viral URI
 - *Staphylococcus aureus*
 - *Haemophilus influenza*
 - Streptococci and pneumococci
- Pathophysiology:
 - Swelling of tracheal mucosa below vocal cords
 - Thick, purulent secretions may lead to mucous plugging

Bacterial Tracheitis

- Presentation similar to croup
 - More toxic appearing child
 - Does not respond well to typical croup treatment
 - Outside the typical age group for croup
- Soft tissue neck film
 - Edema with an irregular border of the subglottic tracheal mucosa
 - “Subglottic membrane”

Bacterial Tracheitis



Bacterial Tracheitis

- Management
 - Assess and maintain patent airway
 - Frequent suctioning if intubated
 - ENT consultation
 - Broad spectrum antibiotic coverage

Foreign Body Aspiration

- Consider this when:
 - Child has recurrent wheezing or stridor unresponsive to typical therapy
 - Afebrile
 - Recurrent pneumonia in same location
- Symptoms may be:
 - Acute – large item in large airway
 - Chronic – small item in small airway; asymptomatic period common


Foreign Body Aspiration

- Common items found:
 - Coins
 - Nuts or seeds
 - Popcorn, small candy
 - Beads, buttons, safety pins
 - Balloons, latex gloves
 - Toys with small or loose parts

Foreign Body Aspiration

- Diagnosis:
 - Soft tissue films of neck
 - PA and lateral chest films
 - Bilateral decubitus films
 - Inspiratory and expiratory chest films
 - Look for air trapping, mediastinal deviation, atelectasis
 - Foreign body itself may be radio-opaque or radio-lucent


Foreign Body Aspiration



Foreign Body Aspiration



Foreign Body Aspiration



Other Upper Airway Problems

- Peritonsillar Abscess
 - Asymmetry of tonsillar pillars
 - Deviation of uvula
- Subglottic Stenosis
 - Common in premature infants that underwent prolonged intubation
- Tracheo/Laryngomalacia
 - Absence of abnormal breath sounds when infant prone


Other Upper Airway Problems

- Neoplasms:
 - Papilloma
 - Vocal cord nodules
- Bronchogenic cysts
- Cystic hygroma (Lymphangioma)
- Vascular rings
- Tracheo-esophageal fistulas
- Laryngeal webs

Papillomas



Tracheo-Esophageal Fistulas



Lower Airway Illnesses

- Intrathoracic Structures
 - Mainstem bronchi, bronchial tree, bronchioles
- Wheezing and Rales
 - Obstruction of intrathoracic airway
 - Heard during expiration and inspiration
 - Air trapping and atelectasis
 - Diminished air movement

Lower Airway Illnesses

- Asthma
- Bronchiolitis
- Bronchopulmonary dysplasia
- Pneumonia

Asthma

- Reversible airway obstruction:
 - Bronchospasm of lower airway
 - Swelling of airways and increased mucous production (inflammation)

Asthma - Triggers

- Atopic conditions
 - allergic rhinitis, eczema, chronic sinusitis
- Allergen exposures
 - Cigarette smoke
 - Pets
 - Carpeting, ceiling fans (dust mites)
 - Cockroaches
- Viral illnesses

Asthma - Presentation

- Cough
- Wheeze
- Shortness of breath
- Chest tightness
- Vomiting
- History:
 - Frequency, duration of symptoms
 - Previous admissions, PICU stays
 - Previous steroid use, varicella exposure


Asthma – Physical Exam

- Assess work of breathing
 - Retractions
 - Nasal flaring
 - Increased respiratory rate
- Assess for hypoxia
- Lung exam
 - Wheezing
 - Prolonged expiratory phase
 - Rhonchi or rales
 - Air movement
 - Absence of wheezing is worrisome

Asthma

- When to order X-ray?
 - Hypoxia
 - Asymmetric lung sounds
 - First time wheezing
- Chest X-ray findings:
 - Hyperinflation
 - Peribronchial cuffing
 - Atelectasis

Asthma – Chest x-ray



Asthma - Management

- Bronchospasm
 - Beta-2 agonists
 - Albuterol – administer via nebulizer with oxygen
 - <10kg: 2.5 mg
 - >10kg: 5.0 mg
 - Anticholinergics
 - Atrovent (peanut allergy a contraindication for atrovent administered by metered dose inhaler, not for nebulized solution)
 - <10kg: 250 mcg
 - >10kg: 500mcg

Asthma - Management

- Inflammation
 - Corticosteroids
 - Prednisone
 - Loading dose 2mg/kg
 - Max dose 60mg
 - Orapred liquid comes in 15mg/5ml solution – tastes great!
 - Solumedrol
 - Loading dose 2mg/kg
 - Max dose 125mg
 - Use when patient vomiting, unable to hold down oral medications
 - Same efficacy as oral steroids

Asthma – Management

- Non-responsive to traditional care:
 - Magnesium Sulfate
 - 25 mg/kg/dose
 - Continuous albuterol
 - 10mg/hour
 - Terbutaline
 - 0.01 mg/kg/dose SQ every 20 minutes x2
 - 0.1-0.4 mcg/kg/minute drip
 - Epinephrine
 - 0.01 mg/kg/dose SQ every 20 minutes x4
 - Consider ketamine as sedative if patient needs intubation

Enterovirus D-68

- New strain of Enterovirus developed this year
 - Most enteroviral infections cause GI symptoms such as vomiting and diarrhea
 - Typically a summer illness
 - This strain was seen later in the season than most enteroviral infections
 - Symptoms include sudden onset asthma-like symptoms in patients with no prior history of asthma
 - Acute onset of wheezing, increased work of breathing
 - Little or no fever
 - Lack of typical GI symptoms
 - Treat like typical asthma patient
 - Symptoms improve rapidly over 48-72 hours

Bronchiolitis

- Similar to asthma but symptoms are caused by viral etiology:
 - RSV
 - Parainfluenza
 - Adenovirus
 - Rhinovirus
 - Mycoplasma

Bronchiolitis

- RSV induces damage to the bronchial epithelium resulting in lower airway inflammation
- Bronchospasm (with history of atopy)
- Most common in winter, early spring
- Age typically < 3 years old
- Symptoms worse in premature infants

Bronchiolitis - Symptoms

- Cough
- Tachypnea
- Accessory muscle use
- High pitched wheezing
- Fine inspiratory crackles; rhonchi
- Copious, thick nasal secretions
- Low grade fever

Bronchiolitis

- Associated findings:
 - Otitis media
 - Pneumonia
 - Apnea (premature infants)
 - Dehydration
- Chest X-ray:
 - Hyperinflation / air trapping
 - Increased perihilar markings
 - Areas of atelectasis

Bronchiolitis - Management

- Mainly supportive
 - Nasal saline spray, frequent suctioning
 - Adequate PO intake (Pedialyte, juice)
- Pharmacologic
 - Albuterol – may diminish wheezing
 - Racemic Epinephrine – may diminish tachypnea
 - Steroids – controversial – consider if history of atopy

Bronchiolitis

- Infants infected with RSV more likely to wheeze with future viral infections due to airway remodeling
- Appropriate Discharge Criteria:
 - No hypoxia
 - Taking PO fluids well
 - No tachypnea or increased work of breathing
 - Wheezing may still be present – consider use of Albuterol MDI with spacer and face mask
 - Reliable parents, follow up


Bronchopulmonary Dysplasia (BPD)

- Chronic lung disease of premature infants that required prolonged respiratory support in NICU:
 - Lung immaturity
 - Oxygen therapy
 - Positive pressure ventilation
 - Infection and inflammation
 - Poor nutrition

BPD

- Symptoms and Findings:
 - Tachypnea, retractions at rest or during mild URI
 - Lungs hyperinflated (increased AP diameter)
 - Crackles, wheezes, decreased breath sounds
 - Chronic CO₂ retention
 - CXR – hyperinflation, cystic areas, atelectasis, coarse appearing

BPD – Chest X-Ray



BPD - Management

- Supportive care
 - Humidified oxygen
 - Frequent suctioning
 - Adequate hydration
 - Assisted ventilation
 - Trial of beta-agonist
- Screen for RSV infection
 - Most infants will receive Synagis

Pneumonia

- Etiology differs in age groups
- Chest X-ray may be useful to help differentiate between different etiologies
- Symptoms and findings:
 - Cough, tachypnea, hypoxia
 - Asymmetric breath sounds, rales, decreased air movement
 - Fever
 - Elevated WBC with left shift

Pneumonia - Neonatal

- Bacterial causes:
 - *E. Coli*
 - Group B Strep
 - *Staph aureus*
 - *Listeria monocytogenes*
- Treatment:
 - Ampicillin and Gentamycin

Pneumonia

- *Chlamydia trachomatis*
 - Consider this in 3 week old to 4 month old patients
 - Afebrile pneumonitis with congestion, wheezing, fine diffuse crackles
 - Paroxysmal cough
 - Prior concomitant inclusion conjunctivitis
 - Treat with erythromycin


Pneumonia

- *Bordetella pertussis*
 - Severe paroxysmal coughing episodes followed by cyanosis and apnea
 - Sometimes associated with inspiratory “whoop” in older children
 - Ask about immunization history
 - CXR: hyperinflation, perihilar infiltrates, atelectasis, or normal
 - CBC: elevated WBC with lymphocytic predominance
 - Send Bordetella FA and culture (call lab for kit)
 - Treat with Erythromycin

Pneumonia

- Infant / child <4 years
 - Lobar – *S. pneumonia*
 - Amoxicillin – no hypoxia; well-hydrated
 - Rocephin IV
 - Atypical
 - Respiratory viruses
 - Influenza

Lobar Pneumonia - Bacterial



Pneumonia

- Older children
 - Lobar
 - *S. pneumonia*
 - Atypical
 - *Mycoplasma pneumoniae*
 - *Chlamydia pneumoniae*
 - Influenza
- Treat with Amoxicillin and/or Zithromax (outpatient)
- Treat with Rocephin and Zithromax (inpatient)

Questions? Comments?

Chris Woleben, M.D.

804-592-8276

cwoleben@vcu.edu