# Virginia Department of Health

# Office of Emergency Medical Services (OEMS)

# **Quarterly Report on Trauma Incidents**

Q4 2020

Division of Trauma and Critical Care 1041 Technology Park Drive Glen Allen, Virginia 23059 Phone: (804) 888-9100

This report is based on the deliberations of the System Improvement Committee and analyses performed by Office of EMS Epidemiology staff.

#### Introduction

Section B 3. of the Code of Virginia (§32.1-111.3) requires the monitoring of the quality of the Commonwealth's emergency medical services (EMS) and trauma services using data from the EMS patient care information system. The EMS Advisory Board reviews and analyzes such data quarterly and reports its findings to the Commissioner. The Advisory Board has delegated this function to the System Improvement Committee (formerly the Trauma Performance Improvement Committee).

This quarterly report focuses on four key areas:

- 1. Completeness of prehospital vital sign documentation (blood pressure, respiratory rate, and Glasgow Coma Score) as required in Step 1 of the Virginia Field Trauma Triage Decision Scheme.
- 2. The number of trauma patients treated and transported by EMS agencies.
- 3. The number of trauma patients who met Step 1 (vitals), Step 2 (anatomy of injury), and Step 3 (mechanism of injury/impact) Virginia Field Trauma Triage Criteria.
- 4. The number of patients meeting trauma triage criteria transported to hospitals not designated as trauma centers.

The results reported here represent a high-level summary of the findings. This report describes how each EMS Council Region is performing. The report will be provided to the appropriate Regional EMS Council Director for each region. The Directors will be given an opportunity to provide feedback, which may explain special circumstances for which an exception occurred. The findings of this report and any feedback from the Directors will be used to drive education and improve the Trauma Triage Plan.

EMS patient data is extracted from patient medical records submitted by EMS agencies to the Virginia Pre-Hospital Information Bridge (VPHIB) program (Elite v3) maintained within the Virginia Department of Health's (VDH) Office of Emergency Medical Services (OEMS) Division of Trauma/Critical Care. Data summarized in this report represent EMS responses that occurred during the fourth quarter of 2020 (October through November) and were entered into VPHIB v3 as of 4/05/2020. VPHIB v3 data are based on the National EMS Information System (NEMSIS) standards.

This report includes all EMS responses categorized as trauma incidents using the following guidelines (Table 1).

Table 1. Definition of Trauma Patients within VPHIB version 3

## **Type of Service Requested**

### 911 Response (Scene)

## **Incident/Patient Disposition**

## Patient Treated, Transported by this EMS unit


## **Situation Provider Primary Impression (ICD-10-CM)**

- S00-S09 (Injuries to the head)
- S10-S19 (Injuries to the neck)
- S20-S29 (Injuries to the thorax)
- S30-S39 (Injuries to the abdomen, lower back, lumbar spine, pelvis, and external genitals)
- S40-S49 (Injuries to the shoulder and upper arm)
- S50-S59 (Injuries to the elbow and forearm)
- S60-S69 (Injuries to the wrist, hand, and fingers)
- S70-S79 (Injuries to the hip and thigh)
- S80-S89 (Injuries to the knee and lower leg)
- S90-S99 (Injuries to the ankle and foot)
- T07 (Injuries involving multiple body regions)
- T14 (Injury of unspecified body region)
- T20-T25 (Burns and corrosions of external body surfaces, specified by site)
- T26-T28 (Burns and corrosions confined to eye and internal organs)
- T30-T32 (Burns and corrosions of multiple and unspecified body regions)
- T75.0 (Effects of lightning)
- T75.4 (Electrocution) (With 7th digit character modifier of A, B, or C; D through S are excluded)

## Excluding:

- S00 (Superficial injuries of the head)
- S10 (Superficial injuries of the neck)
- S20 (Superficial injuries of the thorax)
- S30 (Superficial injuries of the abdomen, pelvis, lower back and external genitals)
- S40 (Superficial injuries of shoulder and upper arm)
- S50 (Superficial injuries of elbow and forearm)
- S60 (Superficial injuries of wrist, hand, and fingers)
- S70 (Superficial injuries of hip and thigh)
- S80 (Superficial injuries of knee and lower leg)
- S90 (Superficial injuries of ankle, foot, and toes)

Figure 1. Virginia Field Trauma Triage Decision Scheme



## Virginia Trauma Summary, Fourth Quarter, 2020

EMS agencies in Virginia responded to a total of 385,157 EMS calls; of that total, 259,304 (67.3%) patients had a disposition of treated and transported by the unit, 42,483 (11%) had a disposition of canceled, 26,570 (6.9%) patients had a disposition of EMS assist, 4,798 (1.2%) patients had a disposition of treated and transferred care to another unit, 4,360 (1.1%) patients were documented as dead at the scene, and 47,642 (12.4%) patients had some other incident disposition (e.g., patient treated and released AMA, patient treated and transported by private vehicle, etc.). Out of the total EMS calls, 22,379 (5.8%) incidents were classified as trauma incidents in VPHIB. The Northern Virginia EMS Council had the highest number of trauma calls (5,062; 22.6%), followed by the Old Dominion EMS Alliance (4,589; 20.5%). Trauma incident numbers for the quarter, broken down by month and Regional EMS Council, are shown in Figure 2. Tables 2-4 summarize the body regions most frequently affected by trauma, the top 10 hospitals receiving trauma transports, and vital signs data quality for trauma incidents.




Figure 2. Monthly Trauma Incidents by Regional EMS Council, Fourth Quarter 2020, Virginia

Table 2. Trauma Incidents by Abbreviated Injury Scale (AIS) Body Region, Fourth Quarter 2020, Virginia

Body Region	Counts of Incidents
Head	4,181 (18.7%)
Face	1,552 (6.9%)
Neck	900 (4.0%)
Thorax	551 (2.5%)
Abdomen	516 (2.3%)
Spine	1,441 (6.4%)
Upper Extremity	3,274 (14.6%)
Lower Extremity	5,348 (23.9%)
Unspecified	4,578 (20.5%)
Multiple Injuries	38 (0.2%)

Table 3: Top Ten Hospital Destinations for Trauma Incidents, Fourth Quarter 2020, Virginia

Destination Hospital For Trauma Incidents	Counts of Incidents
Fairfax Hospital	1,190 (5.3%)
Roanoke Memorial Hospital	920 (4.1%)
Norfolk General Hospital	813 (3.6%)
VCU Health Systems	785 (3.5%)
Chippenham Hospital	745 (3.3%)
Riverside Regional Medical Center	721 (3.2%)
UVA Health System	654 (2.9%)
Northern Virginia Medical Center	610 (2.7%)
Mary Washington Hospital	595 (2.7%)
Virginia Beach General Hospital	521 (2.3%)

Table 4. Vital Signs Data Quality for Trauma Incidents, Fourth Quarter 2020, Virginia

Vital Signs Data Quality	Counts of Incidents
Total Number of Trauma Incidents	22,379
Patients with All 3 Vital Signs Reported	21,813 (97.5%)
Patients with Incomplete* Vital Signs	566 (2.5%)
Patients with Systolic Blood Pressure Reported	22,279 (99.6%)
Patients with Respiratory Rate Reported	22,109 (98.8%)
Patients with Glasgow Coma Score Reported	22,090 (98.7%)

<sup>\*</sup>Incomplete vital signs are missing one or more of the vital signs required in Step 1 of the Trauma Triage algorithm (i.e., Systolic Blood Pressure, Respiratory Rate, or Glasgow Coma Score).

## **Trauma Incidents Meeting Virginia Trauma Triage Criteria**

- Of the 22,379 trauma incidents reported by EMS during the fourth quarter of 2020, 1,814 (8.1%) met Trauma Triage Step 1 criteria, 511 (2.3%) met Step 2 criteria, and 18 (0.1%) met Step 3 criteria. Incidents can meet criteria for more than one step; those incidents were classified into the highest severity level met. For example, if an incident met both Step 1 and Step 2 criteria, it was counted as a Step 1 incident.
  - o Among the incidents meeting Step 1 criteria, 1,493 (82.3%) were classified as meeting Step 1 based on reported vital signs (see Appendix 1). The remaining 321 (17.7%) incidents were classified as meeting Step 1 based on the provider's impression, as reported in the "Trauma Triage Criteria" field in the patient care report.
  - Incidents meeting Step 2 and Step 3 were based solely on the "Trauma Triage Criteria" field.

## Pediatric Patients (Age < 15)

Trauma patients <15 years old are considered pediatric patients per trauma triage criteria. Of the 22,379 trauma incidents reported by EMS during the fourth quarter of 2020, 943 (4.2%) occurred among pediatric patients. Of the 1,814 Virginia trauma incidents meeting Step 1 trauma criteria, 152 (8.4%) occurred among pediatric patients (further details are shown below).

Table 5. Hospital Destination Type for Pediatric Patients Meeting Step 1 Criteria by Regional EMS Council, Fourth Quarter 2020, Virginia

		Trauma Hospital				
Regional EMS Council	Met Step 1	Level I	Level II	Level III	Pediatric Trauma Center	Non-Trauma Hospital
Blue Ridge	4	1	2	0	0	1
Central Shenandoah	9	0	0	0	0	9
Lord Fairfax	6	0	3	1	0	2
Northern	33	12	1	4	1	15
Old Dominion	36	6	2	2	15	11
Peninsulas	5	0	2	0	1	2
Rappahannock	9	0	7	0	0	2
Southwest	9	1	0	2	0	6
Thomas Jefferson	8	8	0	0	0	0
Tidewater	22	2	0	1	14	5
Western	11	0	0	0	8	3
<b>Grand Total</b>	152	30 (19.7%)	17 (11.2%)	10 (6.6%)	39 (25.7%)	56 (36.8%)

- There were 66 incidents involving pediatric patients that met Step 1 trauma criteria that were taken to a Level III trauma center or lower designation. OEMS queried the Virginia State Trauma Registry data to locate those patients and were able to match 5 (7.6%) patients. The deterministic match was performed using the patient's first name, last name, transfer of care date, emergency department (ED) admission date, incident destination hospital, and date of birth as the matching variables. For the 5 patients who matched, patient dispositions included the following:
  - O Two (40%) patients were transferred to another hospital.
  - O Two (40%) patients died at the ED of the hospital.
  - One (20%) patient was taken to the operating room and was later discharged.
- There were 21 pediatric patients who met Step 2 triage criteria. Ten (47.6%) were taken to a pediatric trauma center, 3 (14.3%) were taken to a Level I trauma center, 3 (14.3%) were taken to a Level II trauma center, and 5 (23.8%) were taken to non-trauma designated hospitals.
- No pediatric patients met Step 3 triage criteria.
- There were 98 pediatric patients who received a medication other than oxygen. Of those, 64
  (65.3%) patients had a weight recorded; all weights documented were recorded in kilograms.

## **Geriatric Patients (Age ≥ 65)**

There were 9,547 (42.7% of total trauma incidents) reports of trauma among geriatric patients during the fourth quarter of 2020. Of the 1,814 Virginia trauma incidents meeting Step 1 trauma criteria, 708 (39%) occurred among geriatric patients (further details are shown below).

Table 6. Hospital Destination Type for Geriatric Patients Meeting Step 1 Criteria by Regional EMS Council. Fourth Quarter 2020. Virginia

		Trauma Hospital			Non-Trauma
Regional EMS Council	Met Step 1	Level I	Level II	Level III	Hospital
Blue Ridge	26	3	22	0	1
Central Shenandoah	22	1	0	0	21
Lord Fairfax	15	0	8	0	7
Northern	138	38	8	9	83
Old Dominion	171	64	21	14	72
Peninsulas	64	5	22	0	37
Rappahannock	36	0	23	0	13
Southwest	61	3	7	1	50
Thomas Jefferson	30	22	1	0	7
Tidewater	57	10	2	18	27
Western	88	36	0	8	44
<b>Grand Total</b>	708	182 (25.7%)	114 (16.1%)	50 (7.1%)	362 (51.1%)

- There were 412 incidents involving geriatric patients who met Step 1 trauma criteria who were taken to a Level III trauma center or lower designation. OEMS queried the Virginia State Trauma Registry data to locate those patients and were able to match 67 (16.3%) patients. The deterministic match was performed using the patient's first name, last name, transfer of care date, ED admission date, incident destination hospital, and date of birth as the matching variables. For the 67 patients that matched, patient dispositions included the following:
  - Thirty-three (49.3%) patients were admitted to the hospital; of those, 30 patients were later discharged, one patient was transferred to another hospital, one patient was sent to inpatient rehabilitation, and one patient died at the hospital.
  - Sixteen (23.9%) patients were transferred to another hospital.
  - Six (9%) patients were admitted to the intensive care unit; of those, four were later discharged from the hospital and two died at the hospital.
  - Six (9%) patients were admitted to telemetry and were later discharged.
  - Two (3%) patients were kept in an observation unit; of those, one was later discharged and one was sent to inpatient rehabilitation.
  - o Two (3%) patients died at the ED of the hospital.
  - o Two (3%) patients were discharged from the ED of the hospital.
- Of the 362 geriatric patients who met Step 1 criteria and were taken to non-trauma designated hospitals, 73 (20.2%) had an EMS provider primary impression of an isolated hip injury.
- There were 110 geriatric patients who met Step 2 trauma triage criteria. Of those, 35 (31.8%) patients were taken to a Level I trauma center, 10 (9.1%) were taken to a Level II trauma center, 9 (8.2%) were taken to a Level III trauma center, and 56 (50.9%) were taken to non-trauma designated hospitals.
- Four geriatric patient met Step 3 trauma triage criteria and all the patients were taken to a Level I trauma center.
- For 59 incidents, patient age was recorded to be greater than 100. Quality assurance of a 30% random sample of these incidents showed that 5.6% of the entered ages were incorrect.

## Adult Patients (15 ≥ Age < 65)

The majority of trauma cases that occurred during the fourth quarter of 2020 were among adult patients (n=11,882; 53.1% of all trauma incidents). Of the 1,814 Virginia trauma incidents meeting Step 1 trauma criteria, 951 (52.4%) occurred among adult patients. The hospital destination type for adult trauma incidents meeting Step 1 criteria is shown below by Regional EMS Council (Table 7).

Table 7. Hospital Destination Type for Adult Patients Meeting Step 1 Criteria by Regional EMS Council, Fourth Quarter 2020, Virginia

		Trauma Hospital			Non-Trauma
Regional EMS Council	Met Step 1	Level I	Level II	Level III	Hospital
Blue Ridge	20	8	10	0	2
Central Shenandoah	31	4	0	0	27
Lord Fairfax	21	3	15	0	3
Northern	212	95	22	8	87
Old Dominion	211	133	18	13	47
Peninsulas	59	7	38	0	14
Rappahannock	44	2	27	0	15
Southwest	61	6	14	4	37
Thomas Jefferson	41	28	4	0	9
Tidewater	135	67	4	35	29
Western	113	65	1	9	38
Out of State	3	3	0	0	0
<b>Grand Total</b>	951	421 (44.3%)	153 (16.1%)	69 (7.3%)	308 (32.4%)

- There were 377 incidents involving adult patients who met Step 1 trauma criteria who were taken to a Level III trauma center or lower designation. OEMS queried the Virginia State Trauma Registry data to locate those patients and were able to match 57 (15.1%) patients. The deterministic match was performed using the patient's first name, last name, transfer of care date, incident destination hospital, ED admission date, and date of birth as the matching variables. For the 57 patients that matched, patient dispositions included the following:
  - Eighteen (31.6%) patients were transferred to another hospital.
  - Twelve (21.1%) patients were admitted to the hospital; of those, nine were later discharged from the hospital, two died at the hospital, and one was transferred to another hospital.
  - Eleven (19.3%) patients were admitted to the intensive care unit; of those, seven patients were later discharged, two patients were sent to inpatient rehabilitation, one patient left against medical advice (AMA), and one patient died at the hospital.
  - o Five (8.8%) patients died at the ED of the hospital.
  - o Four (7%) patients were taken to the operating room and were later discharged.
  - Three (5.3%) patients were kept in an observation unit; of those, two were later discharged and one was sent to inpatient rehabilitation.
  - Two (3.5%) patients were admitted to telemetry; of those, one was later discharged and one was sent to inpatient rehabilitation.
  - One (1.8%) patient was discharged from the ED of the hospital.
  - One (1.8%) patient had a ED disposition of 'Other (jail, institution, etc).'

- There were 379 adult patients who met Step 2 criteria. Of those, 243 (64.1%) patients were taken to a Level I trauma center, 42 (11.1%) patients were taken to a Level II trauma center, 33 (8.7%) were taken to a Level III trauma center, and 61 (16.1%) patients were taken to non-trauma designated hospitals.
- There were 14 adult patients who met Step 3 criteria. Of those, 13 (92.9%) were taken to a Level I trauma center and 1 (7.1%) was taken to a non-trauma designated hospital.

## **Air-Medical EMS Transport**

There were 402 trauma patient transports by an air-medical ambulance during the fourth quarter of 2020. Of those:

- Twenty (5%) were pediatric transports, of which:
  - O Seven were taken to a pediatric trauma center, 10 were taken to a Level I trauma center, and 3 was taken to a non-trauma designated hospital.
- Eighty-four (20.9%) were geriatric transports, of which:
  - O Sixty-one were taken to a Level I trauma center, 18 were taken to a Level II trauma center, and 5 were taken to a non-trauma designated hospital.
- Two-hundred and ninty-seven (73.9%) were adult transports, of which:
  - O Two-hundred and fifty-six were taken to a Level I trauma center, 34 were taken to a Level II trauma center, and 7 were taken to a non-trauma designated hospital.

## **Causes of Injury**

Trauma patient records were analyzed to identify the causes of injuries in the Commonwealth of Virginia. Fall injuries occurred most commonly, followed by motor vehicle collision injuries. Causes of injury for the third quarter of 2020 are shown in Table 8.

Table 8. Frequencies and Percentages of Causes of Injury, Fourth Quarter 2020, Virginia

Cause of Injury	Frequency	Percentage of the Total
Falls, slips/trips	9,032	40.4%
MVC-related	4,541	20.3%
Blunt force trauma	886	4.0%
Penetrating trauma	535	2.4%
Firearm	310	1.4%
Non-motorized transport	210	0.9%
Animal-related	133	0.6%
Machine-related	108	0.5%
Self-harm	73	0.3%
Burn, smoke inhalation,		
electrocution, explosion	71	0.3%
Abuse	29	0.1%
Recreational	27	0.1%
Poisoning	21	0.1%
Asphyxiation	13	0.1%
Human bite	7	<0.1%

Table 8. Frequencies and Percentages of Causes of Injury, Fourth Quarter 2020, Virginia (continued)

Cause of Injury	Frequency	Percentage of the Total
Environment/weather-related	4	<0.1%
Overexertion/strain	3	<0.1%
Aircraft	3	<0.1%
Drowning	1	<0.1%
Unspecified	6,372	28.5%
Grand Total	22,379	100.0%

## **Under-Triage of Trauma Incidents Heat Map**

A trauma incident is considered to be under-triaged if the incident met Step 1 or Step 2 trauma triage criteria and the patient was taken to either a Level III trauma center or a non-trauma designated hospital. Injuries to the head, legs, and arms occurred most often among the under-triaged incidents (Table 9).

Table 9. Frequencies and Percentages of Under-Triaged Trauma Patients by AIS Body Region of Injury, Fourth Quarter 2020, Virginia

AIS Region	Frequency	Percentage among Under-Triaged Patients
Head	260	25.5%
Unspecified	247	24.2%
Lower Extremities	222	21.8%
Upper Extremities	112	11.0%
Face	70	6.9%
Spine	34	3.3%
Abdomen	29	2.8%
Thorax	24	2.4%
Neck	21	2.1%
Grand Total	1,019	100%

A heat map of under-triaged trauma incidents based on the incident zip code is shown in Figure 3. The heat map shows where the problem is most prominent and serves as a resource planning tool.




Figure 3. Heat Map Showing the Extent of Under-Triage of Trauma Patients, Fourth Quarter 2020, Virginia

## **Distribution of Trauma Facilities across Virginia**

Trauma centers across Virginia are not uniformly distributed. The upper part of the Northern Virginia EMS Council and parts of Central Virginia (e.g., the greater Richmond area) have greater access to trauma centers, as multiple trauma centers are located within close proximity. Most parts of the Old Dominion EMS Alliance, Central Shenandoah EMS Council, and Western Virginia EMS Council have very limited access to trauma centers. The Central Shenandoah EMS Council and Southwest Virginia EMS Council have no trauma centers within their EMS regions, but are reasonably close to Level II trauma centers in other EMS regions or states. The distribution of trauma centers across Virginia, surrounded by rings showing the geographical areas within a 30-minute drive of each trauma center, is shown below (Figure 4). This map displays which parts of Virginia have limited access to a trauma center.



Figure 4. Trauma Centers across Virginia, Surrounded by 30-Minute Drive Time Rings

## **Data Quality**

Virginia EMS agencies have been working very hard to make sure they provide optimal care to their patients while also making efforts to improve data quality. Over the past years, there has been a significant improvement in trauma triage data quality. Continuation of this improvement is what the System Improvement Committee expects. The OEMS conducted quality assurance checks on trauma triage records from the fourth quarter of 2020, as described below.

- Blank Vital Signs: There were a total of 100 incidents without systolic blood pressure
  documented, 270 incidents without respiratory rate documented, and 289 incidents without
  GCS documented. Ten percent of the incidents from each category were randomly selected for
  further review; the findings are listed below.
  - o In some instances, the time vital signs were taken was not documented. This may occur when vital signs are taken before EMS arrival or when the time is not entered into the record by an EMS provider. When this occurs, Elite v3 is not able to identify any initial vital signs contained in the record. As a result, initial vital signs are identified as missing, even when they are recorded. To improve this data point, the time vital signs are taken should be recorded for every instance, when possible.
  - In some cases, vitals are unable to be obtained due to patient refusal or because the
    patient is a child. Such cases should be documented as Pertinent Negatives (e.g.,
    "Refused" or "Unable to Complete"). Leaving the vital sign field blank and reporting such
    cases only in the patient care narrative will result in the vital sign being identified as
    missing.
  - In some cases, vitals were documented in the patient care narratives. Leaving the vital sign filed blank and reporting the vitals in the patient care narrative will result in the vital sign being identified as missing.

- Of the records sampled, 10% of records missing systolic blood pressure, 96.3% of records missing respiratory rate, and 82.8% of records missing GCS were found to have blank data points because the vital signs were not recorded anywhere in the patient record (i.e., the vital sign field or the patient care narrative).
- Atypical Vital Signs: Atypical vital signs are vitals with extreme values. The cutoff values for vitals
  to be considered atypical are chosen arbitrarily only for quality check and validation purposes.
  For this report, systolic blood pressures with values of less than 40 or greater than 250 and
  respiratory rates of less than 3 or greater than 100 were deemed extreme values. There were 59
  instances of extreme values. Thirty percent of the incidents were randomly selected for further
  review.
  - Among reviewed incidents with extreme values, 50% had the same values captured in the narrative and are therefore considered to be valid.
  - Another 27.8% of the vital signs were not supported by the narrative and are considered to be incorrect.
  - In 22.2% of the cases, vitals were documented as such when the providers were
    unsuccessful in obtaining the vitals. For such cases, vital signs should be documented as
    "Unable to Complete." Entering extreme values in the record makes the data less
    accurate and causes the results of any data analysis to be unreliable.
- Not Applicable/Not Recorded/Blank Trauma Triage Criteria: There were 21,029 trauma
  incidents where the "Trauma Triage Criteria" field was reported to be Not Applicable or was Not
  Recorded or Blank. It is understandable that not all trauma incidents meet trauma triage criteria;
  however, some of these records are incorrectly classified or do not report important
  information.
  - Of those incidents, 1,022 (4.9%) had recorded vitals meeting Step 1 trauma triage criteria.
  - Step 2 and Step 3 trauma triage incidents may also be missing trauma triage criteria and therefore may also be incorrectly classified. However, Steps 2 and 3 trauma triage criteria are not based on vital signs, so the exact amount of misclassification cannot be identified.

#### Blank Age

There were 11 incident records where age was left blank; of those incidents, patient age was found in the patient care narratives for 4 patients and records were updated. Of the remaining seven incidents with missing age, three incidents met Step 1 trauma triage criteria and all three patients were taken to Level I Trauma Centers.

#### **Conclusions**

Many factors influence the decision regarding where a patient is transported. As noted above, trauma centers are not equally distributed across Virginia. In some areas (Southwest Virginia and Northern Virginia), out of state trauma center resources are available. Despite having a total of 12 Level I and Level II trauma centers (combined) in Virginia, as well as access to several other similar facilities in surrounding states, large areas of Virginia remain underserved. The variability of resources across Virginia is often compounded by geographic and (especially in the case of Helicopter or Medevac EMS) weather factors. Although a solution to this problem is beyond the scope of this report,

this variability needs to be considered when comparing the outcomes of pre-hospital trauma patients in Virginia.

Missing vital signs data in EMS records continues to be an area of focus for performance improvement efforts. Currently, about one out of every 40 patients (2.5%) have incomplete vital signs data. During the fourth quarter of 2020, 40% of patients who met Step 1 trauma triage criteria and 23.9% of patients who met Step 2 criteria were taken to non-trauma centers. Acknowledging these data, there may be a need to re-examine how trauma triage criteria are being applied in the field, with an eye towards the existing barriers to trauma center access, including the absence of trauma centers in broad swaths of Virginia. Whether the addition of trauma center resources would allow for improved access and care requires further study.

OEMS staff performed quality assurance on trauma triage data from the fourth quarter of 2020. Specifically, the data values that were reviewed included the vital signs used in Step 1 trauma triage criteria designation, atypical vital sign values, and trauma triage criteria fields listed as not applicable, not recorded, or blank. OEMS will continue to perform these data quality checks and will summarize findings for inclusion in future trauma triage reports.

## Appendix 1: Elite v3 Data Dictionary Elements for Trauma Triage Vital Signs and Trauma Triage Criteria

## eVitals.06 - SBP (Systolic Blood Pressure)

#### Definition

The patient's systolic blood pressure.

National Element	Yes	Pertinent Negatives (PN)	Yes
State Element	Yes	NOT Values	Yes
Version 2 Element	E14_04	Is Nillable	Yes
Usage	Required	Recurrence	1:1

### Associated Performance Measure Initiatives

Airway Cardiac Arrest Pediatric STEMI Stroke Trauma

### Attributes

#### NOT Values (NV)

7701001 - Not Applicable 7701003 - Not Recorded

Pertinent Negatives (PN)

8801005 - Exam Finding Not Present 8801019 - Refused 8801023 - Unable to Complete

### Constraints

Data Type minInclusive maxInclusive

integer 500

## Data Element Comment

Required for ACS-Field Triage and other patient scoring systems.

## eVitals.14 - Respiratory Rate

#### Definition

The patient's respiratory rate expressed as a number per minute.

National Element	Yes	Pertinent Negatives (PN)	Yes
State Element	Yes	NOT Values	Yes
Version 2 Element	E14_11	Is Nillable	Yes
Usage	Required	Recurrence	1:1

## Associated Performance Measure Initiatives

Airway Cardiac Arrest Pediatric STEMI Stroke Trauma

### Attributes

### NOT Values (NV)

7701001 - Not Applicable 7701003 - Not Recorded

Pertinent Negatives (PN) 8801005 - Exam Finding Not Present 8801019 - Refused 8801023 - Unable to Complete

## Constraints

Data Type minInclusive maxinclusive

300 integer

## Data Element Comment

# eVitals.23 - Total Glasgow Coma Score

## Definition

The patient's total Glasgow Coma Score.

National Element	No	Pertinent Negatives (PN)	Yes
State Element	Yes	NOT Values	Yes
Version 2 Element	E14_19	Is Nillable	Yes
Usage	Required	Recurrence	1:1

## Associated Performance Measure Initiatives

Airway Cardiac Arrest Pediatric STEMI Stroke Trauma

## Attributes

NOT Values (NV) 7701001 - Not Applicable 7701003 - Not Recorded 7701005 - Not Reporting

Pertinent Negatives (PN) 8801019 - Refused

8801023 - Unable to Complete

## Constraints

Data Type minInclusive maxInclusive

integer

### Data Element Comment

Can be documented or calculated from EVitals.19 (GCS-Eye), EVitals.20 (GCS-Verbal), and EVitals.21 (GCS-Motor).

## elnjury.03 - Trauma Center Criteria

### Definition

Physiologic and Anatomic Field Trauma Triage Criteria (steps 1 and 2) as defined by the Centers for Disease Control.

National Element	Yes	Pertinent Negatives (PN)	No
State Element	Yes	NOT Values	Yes
Version 2 Element		Is Nillable	Yes
Usage	Required	Recurrence	1 : M

### Associated Performance Measure Initiatives

Trauma

## Attributes

# NOT Values (NV)

7701001 - Not Applicable 7701003 - Not Recorded

CorrelationID

Data Type: string minLength: 0 maxLength: 255

## Code List

Description	
Amputation proximal to wrist or ankle	
Crushed, degloved, mangled, or pulseless extremity	
Chest wall instability or deformity (e.g., flail chest)	
Glasgow Coma Score <= 13	
Open or depressed skull fracture	
Paralysis	
Pelvic fractures	
All penetrating injuries to head, neck, torso, and extremities proximal to elbow or knee	
Respiratory Rate <10 or >29 breaths per minute (<20 in infants aged <1 year) or need for ventilatory support	
Systolic Blood Pressure <90 mmHg	
Two or more proximal long-bone fractures	
	Amputation proximal to wrist or ankle Crushed, degloved, mangled, or pulseless extremity Chest wall instability or deformity (e.g., flail chest) Glasgow Coma Score <= 13 Open or depressed skull fracture Paralysis Pelvic fractures All penetrating injuries to head, neck, torso, and extremities proximal to elbow or knee Respiratory Rate <10 or >29 breaths per minute (<20 in infants aged <1 year) or need for ventilatory support Systolic Blood Pressure <90 mmHg

## Data Element Comment

2011 Guidelines for the Field Triage of Injured Patients - value choices for Steps 1 and 2. For falls, one story is equal to 10 feet.

Code 7701001 - Not Applicable should be used when none of the values listed in the code list for element elnjury.03 apply.

## Version 3 Changes Implemented

Added to better evaluate the CDC-ACS 2011 Guidelines for the Field Triage of Injured Patients.

Website: http://www.cdc.gov/FieldTriage/