


## Ehrlichiosis/Anaplasmosis

Agent(s): Bacteria belonging to the family *Anaplasmataceae*. *Ehrlichia chaffeensis* infects monocytes (a type of white blood cell involved with immune function) and causes an illness called human monocytic ehrlichiosis (HME). *E. ewingii* infects granulocytes (a different category of white blood cells) and causes a disease referred to as an *E. ewingii* infection. *Anaplasma phagocytophilum* also infects granulocytes, causing an illness called human granulocytic anaplasmosis (HGA).

Mode of Transmission: Transmitted to humans through the bite of an infected tick. *E. chaffeensis* and *E. ewingii* may infect adult and nymph stage lone star ticks. *Anaplasma phagocytophilum* may infect nymph stage and adult blacklegged ticks (a.k.a., deer ticks). Transmission of these pathogens occurs when an infected tick bites a person and feeds (i.e., remains attached) on that person for a period of more than 24 hours.

Signs/Symptoms: Illness symptoms commonly include fever, headache, body aches, nausea, anorexia, and vomiting. Patients may exhibit signs of thrombocytopenia (low blood platelet count) and leucopenia (low blood white cell count). Severe forms of illness can result in bleeding disorders, difficulty breathing, meningitis/encephalitis, organ damage and death. Persons with weakened immune systems are prone to develop more severe disease.

Prevention: Common practice should include minimizing tick bites by recognizing tick habitats such as humid forest environments with dense undergrowth or heavy leaf litter, and tall weeds and vegetative ground cover along forest margins, tree lines, forest trails and forest clearings. Repellents containing DEET, Picaridin, BioUD, IR3535, or oil of lemon eucalyptus are effective against ticks and should be applied to exposed areas of skin before entering tick habitats. When in tick-prone habitats, light-colored clothing should be worn with pants legs tucked into socks, and permethrin-based repellents should be applied to clothing, socks and shoes. After visiting tick habitats, a person should thoroughly check all body surfaces for ticks and, if found, carefully remove attached ticks as soon as possible.



A total of 131 cases of ehrlichiosis/anaplasmosis were reported in Virginia during 2011. This is a 41% increase from the 93 cases seen in 2010, and a 136% increase from the five-year average of 55.4 cases per year (Figure 13). The considerable increase in reported cases may be due to numerous factors, including increased knowledge of these diseases among healthcare providers, improvements in diagnosis and reporting, and increased tick populations. The increased tick populations result from increased deer populations, particularly in recently developed suburban areas where deer are difficult to control by hunting. Adult lone star ticks and blacklegged ticks both feed primarily on deer blood for reproduction, and deer also serve as a reservoir for *E. chaffeensis*. Among cases reported in 2011, 100 were specified as HME, 21 were specified as HGA, eight were ehrlichiosis/anaplasmosis unspecified, and two were cases of *E. ewingii* infection.

In 2011, ehrlichiosis/anaplasmosis incidence was highest in the 50 to 59 year age group, with 3.8 cases per 100,000 and in the 60 year and older age group, with 3.2 cases per 100,000. Together, these two age groups accounted for 67% of all identified cases. Incidence generally decreased with age, falling to 1.8 per 100,000 in the 40-49 year age group, and down to 1.0 per 100,000 in the <1 year age group. This pattern of age distribution, where infections occur predominantly among those over the age of 50 years, is typical for ehrlichiosis/anaplasmosis observed in other endemic areas of the United States. Race information was not provided for 45% of the reported cases. Among cases where race data were provided, incidence in the white population was almost double that in the black population (1.1 and 0.6 per 100,000, respectively). The rate in males was higher than the rate in females (1.9 and 1.4 per 100,000, respectively).

In 2011, cases were reported from all regions of the state. The highest incidence was seen in the northwest and southwest regions (2.7 and 2.5 cases per 100,000, respectively.) Rates in the remaining regions ranged from 1.1 to 1.6 per 100,000. The majority of cases (52%) had symptom onset in the second quarter, and 37% had symptom onset in the third quarter. The second and third quarters represent the spring and summer months, when ticks are most likely to feed. During 2011, no deaths were attributed to *Ehrlichia* or *Anaplasma* infections.