

Arthritis by Demographics (Race/Ethnicity, Education, Income, Age, and Gender), Virginia, 2012

		Have you ever been told that you have some form of arthritis, rheumatoid arthritis, gout, lupus, or fibromyalgia?							
		Yes				No			
		Sample Size	Weighted Counts	Weighted Percent (%)	CI	Sample Size	Weighted Counts	Weighted Percent (%)	CI
*Virginia	*State Total	2417	1600111	25.5	24.3 -26.6	4938	4685265	74.5	73.4 - 75.7
Race/Ethnicity	Black/Non-Hispanic	352	283450	25.6	22.3 -28.9	691	825085	74.4	71.1 - 77.7
	Hispanic	43	44510	9.2	5.9 - 12.6	241	436856	90.8	87.4 - 94.1
	Other	98	101594	19.1	15.2 -23.1	307	429607	80.9	76.9 - 84.8
	White/Non-Hispanic	1877	1141674	28.1	26.7 -29.5	3607	2924127	71.9	70.5 - 73.3
Education	< H.S.	300	329301	39.3	34.8 -43.8	310	508814	60.7	56.2 - 65.2
	H.S. or G.E.D.	688	445515	27.2	24.9 -29.6	1106	1191775	72.8	70.4 - 75.1
	Some College	659	489282	27.0	24.6 -29.4	1127	1321346	73.0	70.6 - 75.4
	College Graduate	755	328078	16.8	15.5 -18.1	2361	1625954	83.2	81.9 - 84.5
Income	\$15,000 or less	274	212340	37.3	32.5 -42.2	277	356309	62.7	57.8 - 67.5
	\$15,000 to less than \$25,000	400	282163	33.6	29.6 -37.6	503	557898	66.4	62.4 - 70.4
	\$25,000 to less than \$35,000	243	156509	28.0	24.1 -32.0	376	401701	72.0	68.0 - 75.9
	\$35,000 to less than \$50,000	314	196213	26.0	22.8 -29.2	566	558582	74.0	70.8 - 77.2
	\$50,000 or more	774	511301	19.1	17.6 -20.7	2509	2162380	80.9	79.3 - 82.4
Age	Age 18 to 24	---	---	---	---	327	790641	95.4	92.5 - 98.3
	Age 25 to 34	46	68477	6.4	4.3 - 8.4	732	1009530	93.6	91.6 - 95.7
	Age 35 to 44	126	140498	13.0	10.4 -15.5	883	943359	87.0	84.5 - 89.6

	Age 45 to 54	344	334019	28.0	25.0 -31.1	1009	857716	72.0	68.9 - 75.0
	Age 55 to 64	697	443082	43.3	40.4 -46.2	933	580175	56.7	53.8 - 59.6
	Age 65 or older	1189	575995	53.3	50.9 -55.8	1054	503844	46.7	44.2 - 49.1
Gender	Female	1601	918493	28.3	26.7 -29.8	2734	2328279	71.7	70.2 - 73.3
	Male	816	681618	22.4	20.6 -24.2	2204	2356986	77.6	75.8 - 79.4

1. Data source: Virginia Department of Health, Division of Policy and Evaluation, Behavioral Risk Factor Surveillance Survey, 2012. Weighted counts and weighted percents are weighted to population characteristics.
2. "-" replaces estimates when the unweighted sample size for the denominator was < 20 or the CI half width was > 10 for any cell.
3. Responses of don't know/not sure, refused, or missing were removed from the numerator and denominator in all estimates.
4. *** Use caution in interpreting sample sizes less than 50.
5. CI= 95% confidence interval.
6. *** Arthritis diagnoses include: rheumatism, polymyalgia rheumatica, osteoarthritis (not osteoporosis), tendonitis, bursitis, bunion, tennis elbow, carpal tunnel syndrome, tarsal tunnel syndrome, joint infection, Reiter's syndrome, ankylosing spondylitis, spondylosis, rotator cuff syndrome, connective tissue disease, scleroderma, polymyositis, Raynaud's syndrome, vasculitis (giant cell arteritis, Henoch-Schonlein purpura, Wegener's granulomatosis, polyarteritis nodosa)